

Magnetic and electric AdS solutions in string- and M-theory

Aristomenis Donos, Jerome P. Gauntlett and Christiana Pantelidou

*Blackett Laboratory, Imperial College
London, SW7 2AZ, U.K.*

Abstract

The stability properties of a family of magnetic $AdS_3 \times \mathbb{R}^2$ solutions of $D = 5$, $SO(6)$ gauged SUGRA are investigated in more detail. We construct an analogous family of magnetic $AdS_2 \times \mathbb{R}^2$ solutions of $D = 4$, $SO(8)$ gauged SUGRA, including a family of supersymmetric solutions, and also investigate their stability. We construct supersymmetric domain walls that interpolate between AdS_5 and an $AdS_3 \times \mathbb{R}^2$ solution and also between AdS_4 and an $AdS_2 \times \mathbb{R}^2$ solution which provide stable zero temperature ground states for the corresponding dual CFTs. We also construct new families of electric $AdS_2 \times \mathbb{R}^3$ and $AdS_2 \times \mathbb{R}^2$ solutions.

1 Introduction

With a view towards condensed matter applications, there have been several holographic investigations into the behaviour of strongly coupled gauge theories in the presence of magnetic fields, starting with [1–4]. The main focus of this paper will be on $AdS_3 \times \mathbb{R}^2$ and $AdS_2 \times \mathbb{R}^2$ solutions of string or M-theory that are supported by purely magnetic fields in the $\mathbb{R}^2$ directions. Such solutions are of interest because they provide candidate holographic dual descriptions of the IR limit of the zero temperature ground states of field theories in $d = 4$ and $d = 3$, respectively. For this to be the case, it is certainly necessary that the solutions are stable and, in particular, do not contain any modes that violate the AdS_n BF bound. Such stability is guaranteed if the solutions are supersymmetric.

As far as we are aware, the first constructions of such supersymmetric $AdS_3 \times \mathbb{R}^2$ solutions were presented in [5]. The solutions were obtained by uplifting a “magneto-vac” solution of Romans’ $D = 5$ gauged supergravity [6] either on an S^5 , to obtain a solution of type IIB supergravity, or on the general class of M_6 [7, 8] associated with $AdS_5 \times M_6$ solutions of $D = 11$ supergravity dual to $N = 2$ SCFTs in $d = 4$ [5].

Subsequently, a non-supersymmetric magnetic $AdS_3 \times \mathbb{R}^2$ solution of minimal gauged supergravity was shown to arise as the near-horizon geometry of magnetic black brane solutions at zero temperature¹ [12]. These black brane solutions can again be uplifted on an S^5 , to obtain a solution of type IIB supergravity, but also on general X_5 [13] and N_6 [14] associated with $AdS_5 \times X_5$ and $AdS_5 \times N_6$ solutions of type IIB and $D = 11$ supergravity, respectively, dual to $N = 1$ SCFTs in $d = 4$ [15, 16]. One of the conclusions of this paper is that for the special case of uplifting on the S^5 , for which the dual field theory is just $N = 4$ SYM, the $AdS_3 \times \mathbb{R}^2$ solution studied in [12] is not stable and hence does not describe the zero temperature ground state.

More recently, it was found that these supersymmetric and non-supersymmetric magnetic $AdS_3 \times \mathbb{R}^2$ solutions are members of a more general two-parameter family of solutions [17], with the supersymmetric solution of [5, 6] part of a one parameter sub-family, which can be constructed within a $U(1)^3$ truncation of $D = 5$ $SO(6)$ gauged supergravity. An initial investigation into the stability of the non-supersymmetric $AdS_3 \times \mathbb{R}^2$ solutions of [17] was undertaken in [18], within the context of $SO(6)$ gauged supergravity. Here we will re-examine the analysis of [18] finding results which differ in some respects due to a mixing of modes that was overlooked in [18]. In addition, we will show that a large class of the $AdS_3 \times \mathbb{R}^2$ solutions also suffer from

¹Dyonic extensions were studied in [9–11].

a new kind of instability involving neutral scalar fields that are spatially modulated in the $\mathbb{R}^2$ directions similar to [19–22].

Our results, and those of [18], imply that when the solutions are uplifted on S^5 to give type IIB solutions dual to $N = 4$ SYM theory, the parameter space of potentially stable $AdS_3 \times \mathbb{R}^2$ solutions is now very small, but still non-zero (see figures 1 and 3). While we think it is unlikely that there are any further instabilities within $SO(6)$ gauged supergravity, it is still possible that there others within the full KK spectrum. Note that if we consider the one-parameter family of solutions that lie within Romans’ theory, some of the instabilities that we discuss here, but not all, involve fields lying outside of Romans’ theory. This is relevant when we uplift the relevant solutions not an S^5 to $D = 10$, but on the M_6 of [7] to $D = 11$ [5]. Within minimal gauged supergravity, we do not find any instabilities for the unique magnetic $AdS_3 \times \mathbb{R}^2$ solution.

We will also construct a supersymmetric domain wall solution that interpolates between AdS_5 in the UV and a particular supersymmetric $AdS_3 \times \mathbb{R}^2$ solution in the IR. The example we choose lies within Romans’ theory so this can be uplifted both on S^5 to type IIB and also on the class of M_6 of [7] to $D = 11$. This solution, being supersymmetric, should describe the stable ultimate zero temperature ground state of the corresponding $d = 4$ CFTs when they are placed in a magnetic field.

The instabilities that we find for the non-supersymmetric $AdS_3 \times \mathbb{R}^2$ solutions show that these solutions cannot provide the ultimate IR ground states of dual $d = 4$ field theories when held in a magnetic field. Nevertheless, they are still physically interesting. In general, extending the work of [12], we expect to be able to construct finite temperature black hole solutions which at zero temperature interpolate between AdS_5 in the UV² and a given non-supersymmetric $AdS_3 \times \mathbb{R}^2$ solution in the IR. The instability of the latter indicates that there will be a phase transition at finite temperature described by new types of black hole solutions and the instabilities that we discuss suggest the types of modes that will be involved in constructing them. Some specific magnetic $AdS_3 \times \mathbb{R}^2$ solutions have several different kinds of instabilities including spatially modulated instabilities, driven by neutral scalars, and superconducting instabilities driven either by charged scalars, charged vectors³ or combinations thereof. Our results suggests that there is a rich story involving com-

²In general, in addition to the magnetic field, we expect that some of the scalar fields will also give rise to deformations of operators of the UV CFT.

³The possibility of charged vectors producing superconducting instabilities in the presence of magnetic fields has been recently discussed in [23, 24] based on the older work of [25, 26]. This is also reminiscent of “reentrant superconductivity”, reviewed in [27], that is seen in URhGe [28].

peting phases that would be interesting to elucidate. It is also worth emphasising that since the supersymmetric $AdS_3 \times \mathbb{R}^2$ solutions are at the boundary of unstable non-supersymmetric solutions (see figure 3), the corresponding dual ground states must about different phases and hence have the nature of quantum critical points.

We now turn to top down solutions containing AdS_2 factors that are supported by magnetic fields. Such solutions are particularly interesting since they might provide dual descriptions of locally quantum critical points, which have been shown to be associated with interesting non-Fermi liquid behaviour [29–32] (see [33, 34] for the inclusion of magnetic fields). For $D = 5$ it was shown in [17] that the $U(1)^3$ truncation of $SO(6)$ gauged supergravity admits a non-supersymmetric magnetic $AdS_2 \times \mathbb{R}^3$ solution which can be uplifted on an S^5 to type IIB. However, it was subsequently shown in [35] that, within the same truncation, this suffers from an instability involving neutral scalar fields that are spatially modulated in the $\mathbb{R}^3$ directions.

It has long been known that minimal $D = 4$ gauged supergravity admits a non-supersymmetric magnetic $AdS_2 \times \mathbb{R}^2$ solution (the near horizon limit of the standard magnetic AdS-RN black brane solution) and that this can be uplifted to obtain solutions of $D = 10, 11$ supergravity in a variety of ways [15, 16]. The concluding section of [18] briefly discussed how this solution can be generalised to form part of a larger family of magnetic $AdS_2 \times \mathbb{R}^2$ solutions of the $U(1)^4$ truncation of $SO(8)$ gauge supergravity which can be uplifted to $D = 11$ on S^7 . Here we will flesh out these constructions in a little more detail. We will find a two-parameter locus of supersymmetric solutions that includes the specific examples already mentioned in [18], as special cases. We will also show that a large class of the non-supersymmetric solutions suffer from similar instabilities that we find for the magnetic $AdS_3 \times \mathbb{R}^2$ solutions. In particular we explicitly discuss an instability involving neutral scalar fields that are spatially modulated in the $\mathbb{R}^2$ directions and a simple instability involving charged scalars.

We will also construct a supersymmetric domain wall solution that interpolates between AdS_4 in the UV and a representative magnetic $AdS_2 \times \mathbb{R}^2$ solution in the IR^4 . This solution, being supersymmetric, should describe the stable ultimate zero temperature ground state of the corresponding $d = 3$ CFTs when they are placed in a magnetic field. This provides the first top down holographic description of a locally quantum critical point where stability is guaranteed by supersymmetry, and it will be interesting to investigate the behaviour of fermion response functions for

⁴Note that in the context of $N = 2$ $D = 4$ gauged supergravity, related solutions have been discussed in [36–38].

this background.

In the last part of the paper, we will briefly discuss AdS_2 solutions that are supported by electric fields. Indeed an electric-magnetic duality transformation for the $U(1)^4$ truncation of $D = 4$ $SO(8)$ gauged supergravity provides a simple way to obtain a three parameter family of electric $AdS_2 \times \mathbb{R}^2$ solutions. We find that none of them preserve supersymmetry. Furthermore, we also can use the duality transformation to find a domain wall solution, solving first order equations, that interpolates between AdS_4 and a non-supersymmetric $AdS_2 \times \mathbb{R}^2$ solution⁵. We do not investigate instabilities for these solutions here, but we expect that there will be many: for the special case of the electric $AdS_2 \times \mathbb{R}^2$ solution of minimal gauged supergravity see [40].

In addition, we also construct a two parameter family of electric $AdS_2 \times \mathbb{R}^3$ solutions of the $U(1)^3$ truncation of $D = 5$ $SO(6)$ gauged supergravity. These solutions generalise the solutions of Romans' theory given in [6]. Once again, none of these solutions are supersymmetric. We will again leave a detailed analysis of instabilities to future work, but we note that it was already shown in [22] that the electric $AdS_2 \times \mathbb{R}^3$ solutions of Romans' theory suffer from helical p -wave superconducting instabilities.

2 Magnetic $AdS_3 \times \mathbb{R}^2$ solutions

In this section we will review the magnetic $AdS_3 \times \mathbb{R}^2$ solutions of [6, 17, 18]. We will also construct a supersymmetric domain wall solution that interpolates between AdS_5 in the UV and the supersymmetric $AdS_3 \times \mathbb{R}^2$ solution of [6] in the IR.

2.1 $U(1)^3 \subset SO(6)$ gauged supergravity

We start with the $U(1)^3$ truncation of $D = 5$ $SO(6)$ gauged supergravity [41] that keeps two neutral scalar fields ϕ_a . It is convenient to package the two scalars in terms of three constrained scalars X_i via

$$X_1 = e^{-\frac{1}{\sqrt{6}}\phi_1 - \frac{1}{\sqrt{2}}\phi_2}, \quad X_2 = e^{-\frac{1}{\sqrt{6}}\phi_1 + \frac{1}{\sqrt{2}}\phi_2}, \quad X_3 = e^{\frac{2}{\sqrt{6}}\phi_1}, \quad (2.1)$$

⁵In the context of $AdS_4 \times SE_7$ solutions, a non-supersymmetric flow solution that interpolates between AdS_4 in the UV and an electric $AdS_2 \times \mathbb{R}^2$ in the IR was constructed in [39] and, as yet, has not been shown to suffer from any instabilities.

with $X_1 X_2 X_3 = 1$. The Lagrangian is then given by

$$\mathcal{L} = (R - V) * 1 - \frac{1}{2} \sum_{a=1}^2 * d\phi_a \wedge d\phi_a - \frac{1}{2} \sum_{i=1}^3 (X_i)^{-2} * F^i \wedge F^i + F^1 \wedge F^2 \wedge A^3, \quad (2.2)$$

where

$$V = -4 \sum_{i=1}^3 (X_i)^{-1}. \quad (2.3)$$

Any solution of this theory can be uplifted on an S^5 to obtain an exact solution of type IIB supergravity using the formulae in [41].

This theory can be further truncated to obtain a sector of Romans' $SU(2) \times U(1)$ gauge supergravity theory [6]. This is significant because any solution of Romans' theory can also be uplifted to $D = 11$ supergravity using the general class of M_6 [7] associated with supersymmetric $AdS_5 \times M_6$ solutions of $D = 11$ supergravity that are dual to $N = 2$ SCFTs in $d = 4$ [5]. Specifically, if we set $X_1 = X_2 \rightarrow X$, $F^1 = F^2 \rightarrow F^{(3)}/\sqrt{2}$, $F^3 \rightarrow -G$ we obtain Romans' theory as in [5], after setting the two-form to zero and identifying $F^{(3)}$ with one of the $SU(2)$ gauge-fields. There are two other ways of obtaining Romans' theory: one by setting $X_2 = X_3$, $F^2 = F^3$ and another by setting $X_1 = X_3$, $F^1 = F^3$.

By setting the scalars to zero, $X_1 = X_2 = X_3 = 1$, and also $F^1 = F^2 = F^3$ we obtain minimal $D = 5$ gauged supergravity. Recall that any solution of this theory can be uplifted to type IIB supergravity using the general class of X_5 [13] associated with supersymmetric $AdS_5 \times X_5$ solutions of type IIB supergravity that are dual to $N = 1$ SCFTs in $d = 4$ [15] or to $D = 11$ using the general class of N_6 [14] associated with supersymmetric $AdS_5 \times N_6$ solutions of $D = 11$ supergravity also dual to $N = 1$ SCFTs in $d = 4$ [16].

2.2 The $AdS_3 \times \mathbb{R}^2$ solutions

We now consider the family of magnetic $AdS_3 \times \mathbb{R}^2$ solutions to the equations of motion for (2.2) found in [17], and studied further in [18], given by

$$\begin{aligned} ds_5^2 &= L^2 ds^2(AdS_3) + dx_1^2 + dx_2^2, \\ F^i &= 2q^i dx_1 \wedge dx_2, \quad \phi_1 = f_1, \quad \phi_2 = f_2, \end{aligned} \quad (2.4)$$

where f_a are constants,

$$L^{-2} = \sum_{I=1}^3 (\bar{X}_I)^{-1}, \quad (q^i)^2 = \bar{X}_i, \quad (2.5)$$




Figure 1: The moduli space of magnetic $AdS_3 \times \mathbb{R}^2$ solutions. Any point in the (f_1, f_2) plane, combined with a set of signs for the q^i , gives rise to an $AdS_3 \times \mathbb{R}^2$ solution. The red lines correspond to the locus of solutions that can preserve supersymmetry, for particular choices of the signs. The dashed lines correspond to solutions that can be embedded into Romans' theory and the origin corresponds to solutions that can be embedded in minimal gauged supergravity.

and $\bar{X}_i$ are the on-shell values

$$\bar{X}_1 = e^{-\frac{1}{\sqrt{6}}f_1 - \frac{1}{\sqrt{2}}f_2}, \quad \bar{X}_2 = e^{-\frac{1}{\sqrt{6}}f_1 + \frac{1}{\sqrt{2}}f_2}, \quad \bar{X}_3 = e^{\frac{2}{\sqrt{6}}f_1}. \quad (2.6)$$

The q^i can be chosen to have either sign. Notice that when $f_2 = 0$, for example, these are solutions to Romans' theory, and actually were already presented in [6] and uplifted to $D = 10, 11$ supergravity in [5]. When $f_1 = f_2 = 0$ they are solutions of minimal gauged supergravity.

The supersymmetry of these solutions was analysed in [18] where it was shown that the sum of the q^i , with suitable signs, must vanish. We will review this analysis in the next subsection where we also show that the locus of supersymmetric solutions is given by

$$2 \sum_i \bar{X}_i^2 = \left(\sum_i \bar{X}_i \right)^2. \quad (2.7)$$

We have summarised the moduli space of solutions in figure 1.

The AdS_3 solutions, assuming that they are stable, are dual to $d = 2$ CFTs, and the radius, L , is proportional to the central charge. We find that L has a global maximum for the solution with $f_1 = f_2 = 0$. Along the supersymmetric branches, it is a maximum for the three solutions that can be embedded into Romans' theory and then decreases monotonically away from them.

2.3 Supersymmetric AdS_5 to $AdS_3 \times \mathbb{R}^2$ domain wall

It was already shown in [18] that the magnetic $AdS_3 \times \mathbb{R}^2$ solutions (2.4),(2.5) can be supersymmetric, preserving two Poincaré supersymmetries (i.e. $(0, 2)$ in $d = 2$). Here we would like to show that there are supersymmetric domain wall solutions that interpolate between AdS_5 in the UV and $AdS_3 \times \mathbb{R}^2$ in the IR. We thus consider the ansatz

$$\begin{aligned} ds^2 &= e^{2W} (-dt^2 + dy^2) + d\rho^2 + e^{2U} (dx_1^2 + dx_2^2) , \\ F^i &= 2 q^i dx_1 \wedge dx_2 , \\ \phi_a &= \phi_a(\rho) , \end{aligned} \tag{2.8}$$

where W and U are functions of ρ . We will consider the $N = 1$ supersymmetry transformations as given⁶ in [42]

$$\begin{aligned} \delta\psi_\mu &= \nabla_\mu \varepsilon - \frac{i}{2} \sum_i A_\mu^i \varepsilon + \frac{1}{6} \sum_i X_i \gamma_\mu \varepsilon + \frac{i}{24} \sum_i X_i^{-1} [\gamma_\mu^{\nu\rho} - 4\delta_\mu^\nu \gamma^\rho] F_{\nu\rho}^i \varepsilon , \\ \delta\lambda_a &= \left[-\frac{i}{4} \not{\partial} \phi_a + \frac{i}{2} \sum_j \partial_{\phi_a} X_j + \frac{1}{8} \sum_j \partial_{\phi_a} X_j^{-1} F_{\mu\nu}^j \gamma^{\mu\nu} \right] \varepsilon . \end{aligned} \tag{2.9}$$

To preserve these supersymmetries we require that $\sum_i A_\mu^i = 0$ and hence the magnetic charges q^i should satisfy

$$\sum_i q^i = 0 . \tag{2.10}$$

As noted in [18] there are another three $N = 1$ supersymmetries with different sign choices for the gauge fields and hence the charges. We expect these to correspond to the conditions $q_1 + q_2 - q_3 = 0$, $q_1 - q_2 + q_3 = 0$ and $-q_1 + q_2 + q_3 = 0$. As noted in [18], this means that extra supersymmetry can be preserved only if one of the charges is zero: however from (2.5) we see that this is not possible, since $X_i > 0$.

Turning now to the specific ansatz (2.8), choosing q^i to satisfy (2.10) and imposing the projection conditions

$$\gamma_{\hat{r}} \varepsilon = -\varepsilon, \quad \gamma_{\hat{x}_1 \hat{x}_2} \varepsilon = i\alpha \varepsilon, \quad \alpha = \pm 1 , \tag{2.11}$$

⁶Note that we should set their $g = 1$ and identify our X_i with their $X^I = 1/(3X_I)$.

we obtain

$$\begin{aligned}
-W' + \frac{1}{3} \sum_i X_i - \frac{\alpha}{3} e^{-2U} \sum_i X_i^{-1} q^i &= 0, \\
-U' + \frac{1}{3} \sum_i X_i + \frac{2\alpha}{3} e^{-2U} \sum_i X_i^{-1} q^i &= 0, \\
\phi'_a + 2 \sum_j \partial_{\phi_a} X_j + 2\alpha e^{-2U} \sum_j q^j \partial_{\phi_a} X_j^{-1} &= 0, \\
\left[\partial_\rho - \frac{1}{6} \sum_i X_i + \frac{\alpha}{6} e^{-2U} \sum_i q^i X_i^{-1} \right] \varepsilon &= 0.
\end{aligned} \tag{2.12}$$

From the first and the last equation in (2.12) we derive that $\varepsilon = e^{W/2} \eta$, with η a constant spinor satisfying the projection conditions (2.11).

For the supersymmetric $AdS_3 \times \mathbb{R}^2$ solutions, we should set $W = L^{-1} \rho$, $U = 0$ and $X_i = \bar{X}_i$ in (2.11). We then find the conditions

$$L^{-1} = \frac{1}{2} \sum_i \bar{X}_i, \quad -2\alpha q^i = \bar{X}_i (-2\bar{X}_i + \sum_j \bar{X}_j). \tag{2.13}$$

The latter condition combined with (2.10) leads to the condition (2.7), that we mentioned earlier. For the AdS_5 vacuum solution we set $W = U = R^{-1} \rho$, $\phi_a = 0$ and find $R = 1$.

We now show that there exists supersymmetric solutions that interpolate between AdS_5 in the UV and a supersymmetric $AdS_3 \times \mathbb{R}^2$ solution in the IR. We do this for just one representative solution in the IR, namely the one that exists inside Romans' theory. In fact the entire domain wall solution lies within Romans' theory and so we set $\phi_2 = 0$ and consider the supersymmetric solution with $\phi_1 = (2\sqrt{6}/3) \log 2$ and $q^1 = q^2 = -2^{-1/3}$ and $q^3 = 2^{2/3}$ (we have chosen $\alpha = 1$). Within this truncation, the flow equations (2.12) for the non-trivial functions are given by

$$\begin{aligned}
W' - \frac{1}{3} (2e^{-\frac{1}{\sqrt{6}}\phi_1} + e^{\frac{2}{\sqrt{6}}\phi_1}) - \frac{2^{2/3}}{3} e^{-2U} \left(e^{\frac{1}{\sqrt{6}}\phi_1} - e^{-\frac{2}{\sqrt{6}}\phi_1} \right) &= 0, \\
U' - \frac{1}{3} (2e^{-\frac{1}{\sqrt{6}}\phi_1} + e^{\frac{2}{\sqrt{6}}\phi_1}) + \frac{2^{5/3}}{3} e^{-2U} \left(e^{\frac{1}{\sqrt{6}}\phi_1} - e^{-\frac{2}{\sqrt{6}}\phi_1} \right) &= 0, \\
\phi'_1 - \frac{4}{\sqrt{6}} (e^{-\frac{1}{\sqrt{6}}\phi_1} - e^{\frac{2}{\sqrt{6}}\phi_1}) - \frac{2^{5/3}}{\sqrt{6}} e^{-2U} \left(e^{\frac{1}{\sqrt{6}}\phi_1} + 2e^{-\frac{2}{\sqrt{6}}\phi_1} \right) &= 0.
\end{aligned} \tag{2.14}$$

Close to the $AdS_3 \times \mathbb{R}^2$ solution in the far IR the system of equations (2.14) admits




Figure 2: We have plotted U' (blue) W' (purple) and ϕ (green) as functions of ρ for the supersymmetric domain wall solution interpolating between AdS_5 and $AdS_3 \times \mathbb{R}^2$ that exists in Romans' theory.

the expansion

$$\begin{aligned}
W &= w_0 + L^{-1}\rho + \frac{1}{16} \left(-29 + 3\sqrt{33} \right) c_{IR} e^{L^{-1}\delta\rho} + \dots, \\
U &= c_{IR} e^{L^{-1}\delta\rho} + \dots, \\
\phi_1 &= 2\sqrt{\frac{2}{3}} \ln 2 - \sqrt{\frac{3}{2}} \left(-5 + \sqrt{33} \right) c_{IR} e^{L^{-1}\delta\rho} + \dots,
\end{aligned} \tag{2.15}$$

with w_0 and c_{IR} two constants while $L^{-1} = 3/2^{2/3}$ and $\delta = \frac{1}{3}(-1 + \sqrt{33})$. In the far UV we would like to approach the unit radius AdS_5 vacuum. One can easily see that the equations (2.14) admits the expansion

$$\begin{aligned}
U &= \rho + \mathcal{O}(e^{-4\rho}), \\
W &= \rho + \mathcal{O}(e^{-4\rho}), \\
\phi_1 &= 2^{7/6}\sqrt{3}e^{-2\rho}\rho + c_{UV}e^{-2\rho} + \mathcal{O}(e^{-4\rho}),
\end{aligned} \tag{2.16}$$

where c_{UV} is a constant of integration. From the above expansion we see that the scalar ϕ_1 , which is dual to an operator with conformal dimension $\Delta = 2$ in the UV CFT, has both a VEV and a deformation. Using a shooting method we find that there is a solution to (2.14) with boundary conditions (2.15) and (2.16) with $w_0 \approx -0.10$, $c_{IR} \approx 0.31$ and $c_{UV} \approx -1.97$ as we have indicated in figure 2. This is the supersymmetric domain wall solution.

This solution can be uplifted on S^5 to type IIB or on the class of M_6 [7] to $D = 11$ [5]. The solutions then describe the corresponding dual $d = 4$ CFTs deformed by the presence of the magnetic field and also by the operator dual to ϕ_1 . In particular, the uplifted supersymmetric $AdS_3 \times \mathbb{R}^2$ solutions describes the IR ground state at zero temperature.

3 Instabilities of magnetic $AdS_3 \times \mathbb{R}^2$ solutions

In this section we analyse various instabilities of the magnetic $AdS_3 \times \mathbb{R}^2$ solutions given in (2.4),(2.5) within $SO(6)$ gauged supergravity.

3.1 Spatially modulated instabilities of the neutral scalars

Possible instabilities of the two neutral scalars appearing in the $U(1)^3$ truncation (2.2) of $SO(6)$ gauge supergravity were investigated in [18] and none were found. In that analysis only fluctuations independent of the x_1 and x_2 direction were considered. Here we relax this assumption and find that there are spatially modulated modes violating the BF bound hence leading to instabilities, as summarised in figure 3 (a).

3.1.1 Instabilities of the solutions existing in Romans' theory

Let us first discuss the instabilities of the three lines of solutions that are solutions of Romans' theory (the dashed blue lines in figure 1). We should emphasise at the outset that the unstable modes that we find involve fields lying outside of Romans' theory and hence the instability is not relevant when we uplift such solutions on M_6 to $D = 11$ but only when we uplift them on S^5 to type IIB.

To illustrate we will consider the line of $AdS_3 \times \mathbb{R}^2$ solutions with $f_2 = 0$ and $q^1 = q^2$, which arise in Romans' theory. We consider the field perturbation

$$\begin{aligned}\delta A^1 &= a(t, y, \rho) \sin(k x_1) dx_2, & \delta A^2 &= -a(t, y, \rho) \sin(k x_1) dx_2, \\ \delta \phi_2 &= w(t, y, \rho) \cos(k x_1),\end{aligned}\tag{3.1}$$

where a and w are functions of the AdS_3 coordinates and k is a constant. We find that the linearised equations of motion imply that

$$(\square_{AdS_3} - L^2 M^2) \mathbf{v} = 0,\tag{3.2}$$

where $\mathbf{v} = (a, w)$, $\square_{AdS_3}$ is the Laplacian of the unit radius AdS_3 and the mass matrix is given by

$$M^2 = \begin{pmatrix} k^2 & 2\sqrt{2}q^1 k \\ 4\sqrt{2}q^1(\bar{X}_1)^{-2} k & 4(\bar{X}_1)^{-1} + k^2 \end{pmatrix}.\tag{3.3}$$

Notice for $k = 0$ there is no mixing at all as seen in [18]. After diagonalizing the mass matrix we find the two eigenvalues

$$m_{\pm}^2 = \frac{1}{\bar{X}_1} \left(2 + k^2 \bar{X}_1 \pm 2 \sqrt{1 + 4k^2 \bar{X}_1} \right).\tag{3.4}$$

The minimum mass is achieved on the m_-^2 branch when $k_{min} = \pm \frac{1}{2} \sqrt{15} (\bar{X}_1)^{-1/2}$ giving

$$m_{min}^2 = -\frac{9}{4(2 + \bar{X}_1^3)}. \quad (3.5)$$

The AdS_3 BF bound $L^2 m^2 > -1$ is violated for $f_1 > 2\sqrt{\frac{2}{3}} \ln 2$. Note that when $f_1 = 2\sqrt{\frac{2}{3}} \ln 2$, we have $\bar{X}_1 = 2^{-2/3}$ and the solution will satisfy the supersymmetry condition (2.10) provided that q^3 has opposite sign to that of $q^1 = q^2$. In other words, the supersymmetric solutions are located right at the boundary of the set of solutions where the spatially modulated instabilities set in. It is also worth noting that for the supersymmetric solution the static mode that saturates the BF bound, given explicitly by

$$w(\rho) = c_1 e^{-\frac{\rho}{L}} \cos(|k_{min}| x_1), \quad a(\rho) = c_2 e^{-\frac{\rho}{L}} \sin(|k_{min}| x_1), \quad (3.6)$$

where $|k_{min}| = \sqrt{15}/2^{2/3}$ and $c_1^2/c_2^2 = 2^{7/3}3/5$, preserves the supersymmetries of the background. The sign choice of c_1/c_2 depends on the choice of α in the projector (4.11) and we note that for this solution the sign of α is opposite to that of q^1 .

3.1.2 More general analysis

We now consider perturbations about the full two parameter family of $AdS_3 \times \mathbb{R}^2$ solutions (2.4),(2.5). In general we cannot decouple the metric perturbations and so we consider the time independent perturbation

$$\begin{aligned} \delta g_{tt} &= -\delta g_{zz} = L^2 r^2 h_3(r) \cos(k x_1), \\ \delta g_{x_a x_a} &= h_a(r) \cos(k x_1), \quad a = 1, 2, \\ \delta A^i &= a_i(r) \sin(k x_1) dx_2, \quad i = 1, 2, 3, \\ \delta \phi_a &= w_a(r) \cos(k x_1), \quad a = 1, 2, \end{aligned} \quad (3.7)$$

containing eight independent functions, which we take to be functions of the radial coordinate, r , of AdS_3 space when written in Poincaré coordinates (with boundary located at $r \rightarrow \infty$). Expanding the equations of motion of (2.2) around the solutions (2.4) we find a total of eleven differential equations. After a little algebra we can show that for $k \neq 0$ the independent equations for the radial functions consist of two first order equations for h_1 and h_3 and six second order equations for each of the a_i , w_a and h_2 . Note that the equations governing the perturbation (3.7) are independent of the sign choices in (2.5).

To find the scaling dimensions of the dual conformal field theory we look for solutions where the eight functions, as a vector, are of the form $\mathbf{v}r^\delta$ where $\mathbf{v}$ is a constant vector and δ is a constant that is related to a scaling dimension in the two-dimensional conformal field theory dual to the AdS_3 solution. The system of equations then takes the form $\mathbf{M}\mathbf{v} = 0$ where $\mathbf{M}$ is an 8×8 matrix. Demanding that non-trivial values of $\mathbf{v}$ exist implies that $\det \mathbf{M} = 0$ and this specifies the possible values of δ as a function of k . In Figure 3 (a) we have shaded the region of the $f_1 - f_2$ plane for which we find a mode with complex scaling dimension. All the unstable modes that we find are spatially modulated with $k \neq 0$.

It is interesting to note that the boundary of the region plotted in Figure 3 (a) (red curve) is the set of points (f_1, f_2) for which there exists a choice in the signs of (2.5) satisfying the supersymmetry condition. In other words, in the supersymmetric solutions there always exists a mode which saturates the AdS_3 unitarity bound at finite k . Based on the Romans' case, we expect that this mode is always supersymmetric.

At the end of the last section we constructed a supersymmetric domain wall solution interpolating between AdS_5 in the UV and $AdS_3 \times \mathbb{R}^2$ in the IR. These solutions describe the zero temperature ground states of the dual $d = 4$ CFTs when held in a magnetic field and also deformed by the operator dual to the scalar field. The fact that the supersymmetric $AdS_3 \times \mathbb{R}^2$ solutions are at the boundary of unstable $AdS_3 \times \mathbb{R}^2$ solutions indicates that the supersymmetric ground states must adjoin different phases and hence have the nature of quantum critical points.

3.2 Instabilities of some charged scalars

We now want to consider possible instabilities of the magnetic $AdS_3 \times \mathbb{R}^2$ solutions with respect to other fields within $SO(6)$ gauged supergravity. We first recall that $SO(6)$ gauged supergravity has 42 scalar fields, parametrising the coset $E_{6(6)}/USp(8)$ and transforming as $\mathbf{20} + \mathbf{10} + \bar{\mathbf{10}} + \mathbf{1} + \mathbf{1}$ of $SO(6)$. The scalars in the $\mathbf{20}$ irrep are described by a unimodular 6×6 matrix T which, as we will discuss, contains the two scalars in the $U(1)^3$ truncation (2.2) that we have been discussing. In this subsection and the next, we will show that there are additional instabilities⁷ involving the scalar fields in T . Our most general analysis will utilise the consistent truncation [44] that

⁷In appendix A we show that there are no instabilities in a truncation of [43] that keeps four complex scalars in the $\mathbf{10} + \bar{\mathbf{10}}$ irreps (we thank N. Bobev for suggesting this calculation). We also note that the two singlet scalars comprise the axion and dilaton; the stability of the dilaton was discussed in [18]. We have not investigated the stability of these or the 12 two-forms of $SO(6)$ gauged supergravity.




Figure 3: The shaded regions in the (f_1, f_2) plane indicate $AdS_3 \times \mathbb{R}^2$ solutions for which we find have identified one or more unstable modes in $SO(6)$ gauged supergravity. Panel (a) indicates spatially modulated instabilities of neutral scalar fields, discussed in section 3.1, panel (b) indicates instabilities of the charged scalar modes discussed in section 3.2 and panel (c) indicates instabilities of charged vector fields discussed in section 3.3. Not shown are additional instabilities of mixed charged scalars and vectors discussed in section 3.3 that appear in subsets of the blue region. The supersymmetric solutions lie on the red lines.

keeps T and the 15 $SO(6)$ gauge-fields. It is worth noting that when uplifting to type IIB, only the $D = 10$ metric and five-form are involved in this truncation.

As the general analysis using the truncation [44] is rather involved we first investigate possible instabilities of three of the twenty scalars, φ_i , appearing in T , using the consistent truncation of $SO(6)$ gauge supergravity [45]. The Lagrangian is given by

$$\begin{aligned} \mathcal{L} = & (R - V) * 1 - \frac{1}{2} \sum_{a=1}^2 *d\phi_a \wedge d\phi_a - \frac{1}{2} \sum_{i=1}^3 (X^i)^{-2} * F^i \wedge F^i + F^1 \wedge F^2 \wedge A^3 \\ & - \frac{1}{2} \sum_{i=1}^3 *d\varphi_i \wedge d\varphi_i - 2 \sum_{i=1}^3 \sinh^2 \varphi_i * A^i \wedge A^i, \end{aligned} \quad (3.8)$$

with

$$\begin{aligned} V = & -2[2X^2X^3 \cosh \varphi_2 \cosh \varphi_3 + 2X^3X^1 \cosh \varphi_3 \cosh \varphi_1 + 2X^1X^2 \cosh \varphi_1 \cosh \varphi_2 \\ & - (X^1)^2 \sinh^2 \varphi_1 - (X^2)^2 \sinh^2 \varphi_2 - (X^3)^2 \sinh^2 \varphi_3]. \end{aligned} \quad (3.9)$$

After expanding around the background (2.4), the equation of motion for the

charged scalar φ_1 gives

$$\square_{AdS_3}\varphi_1 + L^2\square_{\mathbb{R}^2}\varphi_1 + 4L^2\left(\bar{X}_1\bar{X}_3 + \bar{X}_1\bar{X}_2 - \bar{X}_1^2 - (A^1)^2\right)\varphi_1 = 0, \quad (3.10)$$

where, again, $\square_{AdS_3}$ is the Laplacian of the unit radius AdS_3 . The equations for $\varphi_{2,3}$ are given by cyclic permutations of indices. We now choose a gauge such that $A^i = q^i(x_1 dx_2 - x_2 dx_1)$ while for the scalar we consider the lowest Landau level “ground state”

$$\varphi_1 = e^{-|q_1|(x_1^2+x_2^2)}\psi_1(t, r, z), \quad (3.11)$$

giving

$$(\square_{AdS_3} - L^2 m_{\psi_1}^2)\psi_1 = 0, \quad (3.12)$$

where

$$m_{\psi_1}^2 = -4\left(\bar{X}_1\bar{X}_3 + \bar{X}_1\bar{X}_2 - \bar{X}_1^2 - |q_1|\right), \quad (3.13)$$

which agrees with the last line of equation (6.7) in [18] (after setting their $g = 1$).

In figure 3 (b) we have indicated where these modes violate the BF bound. As one can see from figure 3 (b), these modes intersect the locus of supersymmetric solutions in six places. Using the results of [46] (see eq. (2.24) and set $g = 1$) one can check that at these points the modes saturating the BF bound are supersymmetric, just as we saw in section 3.1.1. The higher Landau levels have mass

$$m_{\psi_1}^2 = -4\left(\bar{X}_1\bar{X}_3 + \bar{X}_1\bar{X}_2 - \bar{X}_1^2 - |q_1|(2n+1)\right), \quad (3.14)$$

again as in [18]. These are unstable in sub-regions of figure 3 (b), and in particular do not intersect the supersymmetric locus.

It is interesting to note that all of these instabilities involve electrically charged fields and hence are associated with new branches of finite temperature superconducting black brane solutions. As the superconductivity is being driven by a magnetic field, it would be interesting to construct and study them further.

3.3 Instabilities of charged scalars and vectors

We will now examine perturbations of $SO(6)$ gauged SUGRA contained in the truncation [44]. This contraction contains twenty scalar fields arranged in a unimodular, 6×6 symmetric matrix T_{ij} and keeps all of the $SO(6)$ gauge fields. The vector and

scalar equation of motion we would like to perturbatively expand are

$$\begin{aligned}
& -D * DT_{ij} + T^{-1k_1k_2} DT_{ik_1} \wedge *DT_{k_2j} = -2 (T_{ik_1} T^{k_1k_2} T_{k_2j} - T_{ik_1} T^{k_1}{}_j T_k{}^k) \\
& + T_{lm}^{-1} * F^l{}_i \wedge F^m{}_j - \frac{1}{6} T_{ij} \left[-2 (T_{lk} T^{lk} - (T_k{}^k)^2) + T_{pk}^{-1} T_{lm}^{-1} * F^{lk} \wedge F^{mp} \right], \\
& D (T_{ik}^{-1} T_{jl}^{-1} * F^{kl}) = -2 T_{k[i}^{-1} * DT_{j]}{}^k,
\end{aligned} \tag{3.15}$$

where

$$\begin{aligned}
DT_{ij} &= dT_{ij} + A_i{}^k T_{kj} - T_{ik} A_j{}^k, \\
F_{ij} &= dA_{ij} + A_{ik} \wedge A_j{}^k,
\end{aligned} \tag{3.16}$$

and we note that the first line of (3.15) corrects a sign in [44].

We find it convenient to switch to a complex notation which just keeps the $SU(3) \subset SO(6)$ symmetry manifest. We will write the magnetic $AdS_3 \times \mathbb{R}^2$ solutions (2.4),(2.5) as

$$\tilde{T}_{I\bar{I}} = X_I, \quad \tilde{A}_{I\bar{I}} = q_I (\bar{z} dz - z d\bar{z}), \tag{3.17}$$

where $I, J = 1, \dots, 3$, are $SU(3)$ indices and $z = \frac{1}{\sqrt{2}} (x^1 + i x^2)$. Note that X_I are the on-shell values, $X_I = \bar{X}_i$, and also $q_I = q^i$ (hence $q_I = \pm X_I^{1/2}$). Consider the scalar perturbation $T = \tilde{T} + t$ where t is a complex matrix. The perturbations $t_{I\bar{I}}$ correspond to perturbations of the neutral scalar fields that we considered in section 3.2, while the perturbations t_{II} correspond to the charged scalars⁸ that we considered in section 3.2. Thus we now consider perturbations t_{IJ} and $t_{I\bar{J}}$ with $I \neq J$. We note that these modes were considered in [18] but the mixing between these modes and the charged modes in the gauge fields was overlooked and we will obtain different results for the spectrum.

We thus consider

$$T = \tilde{T} + t, \quad A = \tilde{A} + a, \tag{3.18}$$

where we are expanding around the background (3.17). Furthermore, we find that it is consistent to set the components of the one-form a along the AdS_3 directions to vanish and so we write

$$a = a^1 dz + a^2 d\bar{z}, \tag{3.19}$$

where a^i and t are complex matrices that are functions of both the AdS_3 coordinates and also $(z, \bar{z})$.

⁸Note that there are three more scalar fields complementing the real φ_i that we didn't explicitly consider.

We will first consider the modes a_{IJ}^i , t_{IJ} , $I \neq J$. Linearising the equations of motion and introducing an appropriate set of ladder operators we are led to the following spectrum (see appendix B for details). Here we define

$$\omega_{IJ} = q_I + q_J, \quad W_{IJ} = \text{sign}(\omega_{IJ})(q_I^{-1} - q_J^{-1}), \quad V_{IJ} = X_I - X_J. \quad (3.20)$$

(note that W is defined slightly differently in the appendix). When $\omega_{IJ} \neq 0$, the independent modes are labelled by two integers $n, m > 0$. There is a tower of modes with $n = 0$, just involving the charged vector fields a_{IJ}^i , which have AdS_3 mass given by

$$m_{0,m}^2 = -2|\omega_{IJ}| - 2W_{IJ}V_{JI} + V_{IJ}^2. \quad (3.21)$$

Also, for each m and n there are mixed modes, involving both a_{IJ}^i and t_{IJ} , with mass matrix

$$M_{n,m}^2 = \begin{pmatrix} 2|\omega_{IJ}|(2n+1) - 4q_I^{-1}q_J^{-1} + 2W_{IJ}V_{JI} + V_{IJ}^2 & 4\sqrt{2|\omega_{IJ}|}W_{IJ}(n+1)^{1/2} \\ 2\sqrt{2|\omega_{IJ}|}W_{IJ}(n+1)^{1/2} & 2|\omega_{IJ}|(2n+1) - 2W_{IJ}V_{JI} + V_{IJ}^2 \end{pmatrix}. \quad (3.22)$$

In figure 3 (c) we have indicated where the zero modes (3.21) can violate the AdS_3 BF bound leading to an instability. The modes arising from diagonalising (3.22) can also violate the BF bound, but only in the region outside of the three supersymmetry lines (in particular there is no overlap with the zero mode instabilities in figure 3 (c)). It is straightforward to determine, numerically, which of the diagonalised modes has the largest violation of the BF bound. The fact that n is an integer leads to a more elaborate structure as compared to the spatially modulated modes labelled by a continuous variable in figure 3 (a).

When $\omega_{IJ} = 0$, which occurs along the three lines with $X_I = X_J$ (the dashed lines in figure 1), we are essentially led back to the mass matrix that we saw for the spatially modulated neutral scalars in (3.3).

The story for the modes $a_{I\bar{J}}^i$, $t_{I\bar{J}}$, with $I \neq J$ is very similar. We now define

$$\omega_{I\bar{J}} = q_I - q_J, \quad W_{I\bar{J}} = \text{sign}(\omega_{I\bar{J}})(q_I^{-1} + q_J^{-1}). \quad (3.23)$$

When $\omega_{I\bar{J}} \neq 0$, the independent modes are again labelled by two integers $n, m > 0$. Again there is a tower of modes with $n = 0$, again just involving the vector fields, with AdS_3 mass given by

$$m_{0,m}^2 = -2|\omega_{I\bar{J}}| - 2W_{I\bar{J}}V_{JI} + V_{I\bar{J}}^2. \quad (3.24)$$

Then for each m and n there are mixed modes with mass matrix

$$M_{n,m}^2 = \begin{pmatrix} 2|\omega_{I\bar{J}}| (2n+1) + 4q_I^{-1} q_J^{-1} + 2W_{I\bar{J}} V_{JI} + V_{IJ}^2 & 4\sqrt{2|\omega_{I\bar{J}}|} W_{IJ} (n+1)^{1/2} \\ 2\sqrt{2|\omega_{I\bar{J}}|} W_{IJ} (n+1)^{1/2} & 2|\omega_{I\bar{J}}| (2n+1) - 2W_{I\bar{J}} V_{JI} + V_{IJ}^2 \end{pmatrix}. \quad (3.25)$$

The instabilities are similar to those we saw above. In particular, the zero modes (3.24) can violate the AdS_3 BF bound leading to an instability as indicated in figure 3 (c)), and there can also be BF violating modes in (3.25). When $\omega_{I\bar{J}} = 0$ the situation is again analogous to the spatially modulated neutral scalars in (3.3).

Note that the instabilities of these charged modes, mixing vectors and scalars, are again associated with holographic superconductivity. These instabilities are somewhat similar to those involving only charged gauge fields in the presence of a magnetic field that were studied in [23], building on [24–26]. We also note that there are various modes which we have studied in this section that saturate the BF bound and intersect with the locus of supersymmetric solutions. We expect them to preserve the supersymmetry but we have not checked the details.

3.4 Discussion

Figure 3 summarises most⁹ of the instabilities that we have found within $SO(6)$ gauge supergravity which are thus relevant to $N = 4$ SYM theory after uplifting on S^5 . We see that apart from the supersymmetric solutions, there is only a very small range of parameters for which we have not found an instability. It would be interesting to know whether or not those solutions are in fact stable within type IIB supergravity. The general picture that has emerged here and in [18] is that studying $N = 4$ SYM in a magnetic field using holography is not a straightforward proposition.

It is worth discussing which of the instabilities that we have discussed reside within the truncation to Romans' theory. This is relevant if we uplift the solutions not on S^5 to type IIB but on the general class M_6 of [7] to $D = 11$ [5]. To be specific we consider the truncation $\phi_2 = 0$, i.e. $X_1 = X_2 \equiv X$ and $A^1 = A^2$. As we already mentioned there is no longer spatially modulated instabilities of the neutral scalar X . In the language of this section, the three $SU(2)$ gauge fields of Romans' theory can be identified with the real $A_{1\bar{1}} = A_{2\bar{2}}$ and the complex A_{12} . Recall that Romans' theory does not have charged scalar fields. Putting this together, we find that instabilities only arise in (3.21) with $IJ = 12$ after noting that $q^1 = q^2$, $W_{12} = V_{12} = 0$. Indeed

⁹It does not include the instabilities arising from (3.22) and (3.25) which, as we discussed, lie in a subset of the cyan coloured regions.

we find instabilities for the solutions in the range $-2.00 \lesssim f_1 \lesssim 0.87$ (recall that the supersymmetric solution has $f_1 \sim 1.13$).

Finally, we note that none of the instabilities that we have discussed appear in minimal gauged supergravity. This is relevant when we uplift the single magnetic $AdS_3 \times \mathbb{R}^2$ solution on the general class X_5 [13] to type IIB supergravity [15] or on the general class of N_6 [14] to $D = 11$ supergravity [16].

4 Magnetic $AdS_2 \times \mathbb{R}^2$ solutions

In this section we construct magnetic $AdS_2 \times \mathbb{R}^2$ solutions of $D = 4$ $SO(8)$ gauged supergravity, which are analogous to the $D = 5$ solutions of section 2. We also construct a supersymmetric domain wall solution that interpolates between AdS_4 in the UV and a supersymmetric $AdS_2 \times \mathbb{R}^2$ solution in the IR.

4.1 $U(1)^4 \subset SO(8)$ gauged supergravity

We consider the $U(1)^4$ truncation of $D = 4$ $SO(8)$ gauged supergravity that keeps three neutral scalar fields ϕ_a [41]. The Lagrangian is given by

$$\mathcal{L} = \frac{1}{2}R - \frac{1}{4} \sum_{a=1}^3 (\partial\phi_a)^2 - \sum_{i=1}^4 X_i^{-2} (F^i)_{\mu\nu} (F^i)^{\mu\nu} - V(X_i), \quad (4.1)$$

where

$$X_1 = e^{\frac{1}{2}(-\phi_1 - \phi_2 - \phi_3)}, \quad X_2 = e^{\frac{1}{2}(-\phi_1 + \phi_2 + \phi_3)}, \quad X_3 = e^{\frac{1}{2}(\phi_1 - \phi_2 + \phi_3)}, \quad X_4 = e^{\frac{1}{2}(\phi_1 + \phi_2 - \phi_3)},$$

$$V(X_i) = -\frac{1}{2} \sum_{i \neq j} X_i X_j = -2(\cosh \phi_1 + \cosh \phi_2 + \cosh \phi_3), \quad (4.2)$$

and we note $X_1 X_2 X_3 X_4 = 1$. Any solution of this theory that satisfies $F^i \wedge F^j = 0$ can be uplifted¹⁰ to $D = 11$ using the formulae in [41]; all of the solutions and the linearised modes that we consider in this section satisfy this condition.

Note that it is consistent to further truncate by setting $X_2 = X_3 = X_4$ along with $F^2 = F^3 = F^4$ to obtain a sector of the $SU(3)$ invariant subsector of $SO(8)$ gauged supergravity [48] [49] and the corresponding uplifted solutions will have $SU(3) \times U(1)^2$ symmetry. This is a case that we will sometimes focus on in the sequel. Alternatively it is also consistent to set $X_1 = X_2$, $X_3 = X_4$ along with $F^1 = F^2$ and $F^3 = F^4$

¹⁰ To do this we should set $g^2 = 1/2$ in eq. (3.8) of [41] and identify $(F^i)^{there} = 2\sqrt{2}(F^i)^{here}$. It is also worth noting that we are using the same conventions as in [47] setting $g = 1$ there.

and the corresponding uplifted solutions will have $SU(2)^2 \times U(1)^2$ symmetry. Both of these theories can be further truncated to minimal gauged supergravity by setting all of the scalars to zero, $X_1 = X_2 = X_3 = X_4 = 1$, and $F^1 = F^2 = F^3 = F^4$. Solutions of minimal $D = 4$ gauged theory can be uplifted to $D = 10$ and $D = 11$ using manifolds associated with general classes of $AdS_4 \times M_7$ solutions dual to $N = 2$ SCFTs in $d = 3$ [16], including SE_7 and also those of section 7.2 of [50].

We now look for the most general class of $AdS_2 \times \mathbb{R}^2$ solutions to the equations of motion of (4.1) that are supported by magnetic fluxes. We thus consider

$$\begin{aligned} ds^2 &= L^2 ds^2(AdS_2) + dx_1^2 + dx_2^2, \\ F^i &= \frac{1}{2} q^i dx_1 \wedge dx_2, \\ \phi_1 &= f_1, \quad \phi_2 = f_2, \quad \phi_3 = f_3, \end{aligned} \tag{4.3}$$

where q^i, f_a are constants and L is the AdS_2 radius. If we define the on-shell quantities

$$\bar{X}_1 = e^{\frac{1}{2}(-f_1-f_2-f_3)}, \quad \bar{X}_2 = e^{\frac{1}{2}(-f_1+f_2+f_3)}, \quad \bar{X}_3 = e^{\frac{1}{2}(f_1-f_2+f_3)}, \quad \bar{X}_4 = e^{\frac{1}{2}(f_1+f_2-f_3)}, \tag{4.4}$$

we find that there is a three parameter family of solutions specified by arbitrary values of (f_1, f_2, f_3) with

$$(q^i)^2 = \frac{\bar{X}_i^2}{2} \sum_{j \neq k \neq i} \bar{X}_j \bar{X}_k, \quad L^{-2} = -2 V(\bar{X}_i), \tag{4.5}$$

and we note that the q^i can be chosen to have either sign.

For the $SU(3) \times U(1)^2$ symmetric subspace of solutions we take $f_1 = f_2 = f_3$ and $q^2 = q^3 = q^4$. This gives rise to a one-dimensional family of solutions labelled by $\bar{X} \equiv \bar{X}_2 = \bar{X}_3 = \bar{X}_4 = (\bar{X}_1)^{-1/3}$ with

$$(q^1)^2 = \frac{3}{\bar{X}^4}, \quad (q^2)^2 = (2 + \bar{X}^4), \quad L^2 = \frac{\bar{X}^2}{6(1 + \bar{X}^4)}. \tag{4.6}$$

4.2 SUSY fixed points

We next want to investigate which of these magnetic $AdS_2 \times \mathbb{R}^2$ solutions are supersymmetric. In the next subsection we will also investigate the possibility of supersymmetric flows that interpolate between AdS_4 in the UV and $AdS_2 \times \mathbb{R}^2$ in the IR. We thus consider the ansatz

$$\begin{aligned} ds^2 &= -e^{2W} dt^2 + d\rho^2 + e^{2U} (dx_1^2 + dx_2^2), \\ F^i &= \frac{1}{2} q^i dx_1 \wedge dx_2, \\ \phi_a &= \phi_a(\rho). \end{aligned} \tag{4.7}$$

where W, U are functions of ρ .

The supersymmetry variations for the $U(1)^4$ truncation of $SO(8)$ gauged supergravity were analysed in [47] and it was shown that it is convenient to break up the $\mathcal{N} = 8$ real Killing spinors into four pairs. We would like to preserve some of the Poincaré supersymmetries of AdS_4 and a consideration of equations (2.15)-(2.16) in [47] implies that depending on which of the four pairs of supersymmetries that we want to preserve we should impose one of the conditions

$$\begin{aligned} q^1 + q^2 + q^3 + q^4 &= 0, & q^1 + q^2 - q^3 - q^4 &= 0, \\ q^1 - q^2 + q^3 - q^4 &= 0, & q^1 - q^2 - q^3 + q^4 &= 0. \end{aligned} \quad (4.8)$$

For definiteness, let us choose to preserve the supersymmetries corresponding to

$$\sum_i q^i = 0. \quad (4.9)$$

The associated supersymmetry variations were written down in (4.3) of [47]. After suitably comparing our notation with that of [47], and switching from two real spinor parameters to a complex spinor parameter ϵ , we have

$$\begin{aligned} \frac{1}{2} \delta \psi_\mu &= \nabla_\mu \epsilon + i \sum_i A_\mu^i \epsilon + \frac{1}{4\sqrt{2}} \sum_i X_i \gamma_\mu \epsilon - i \frac{1}{4\sqrt{2}} \sum_i X_i^{-1} \not{F}^i \gamma_\mu \epsilon, \\ 2 \delta \chi_a &= \left[i\sqrt{2} \not{\partial} \phi_a - i2 \sum_j \partial_{\phi_a} X_j + 2 \sum_j \partial_{\phi_a} X_j^{-1} \not{F}^j \right] \epsilon. \end{aligned} \quad (4.10)$$

Turning now to the specific ansatz (4.7), choosing q^i to satisfy (4.9) and imposing the projection conditions

$$\gamma_{\hat{r}} \epsilon = -\epsilon, \quad \gamma_{\hat{x}_1 \hat{x}_2} \epsilon = i\alpha \epsilon, \quad \alpha = \pm 1, \quad (4.11)$$

we obtain

$$\begin{aligned} -W' + \frac{1}{2\sqrt{2}} \sum_i X_i + \frac{\alpha}{2\sqrt{2}} e^{-2U} \sum_i X_i^{-1} q^i &= 0, \\ -U' + \frac{1}{2\sqrt{2}} \sum_i X_i - \frac{\alpha}{2\sqrt{2}} e^{-2U} \sum_i X_i^{-1} q^i &= 0, \\ -\sqrt{2} \phi'_a - 2 \sum_j \partial_{\phi_a} X_j + 2\alpha e^{-2U} \sum_j q_j \partial_{\phi_a} X_j^{-1} &= 0, \\ \left[\partial_\rho - \frac{1}{4\sqrt{2}} \sum_i X_i - \frac{\alpha}{4\sqrt{2}} e^{-2U} \sum_i q^i X_i^{-1} \right] \epsilon &= 0. \end{aligned} \quad (4.12)$$

From the first and the last equation in (4.12) we derive that $\varepsilon = e^{W/2}\eta$ with η a constant spinor satisfying the projection conditions (4.11).

To determine which of the $AdS_2 \times \mathbb{R}^2$ solutions, summarised in (4.3), (4.4), (4.5), are supersymmetric, we set $W = L^{-1}\rho$, $U = 0$ and $X_i = \bar{X}_i$ in (4.11). We find that the magnetic charges are given by

$$2\alpha q_i = \bar{X}_i(-2\bar{X}_i + \sum_j \bar{X}_j), \quad (4.13)$$

and (4.9) then gives the condition

$$2 \sum_i \bar{X}_i^2 = \left(\sum_i \bar{X}_i \right)^2, \quad (4.14)$$

analogous to what we saw in the $D = 5$ case (2.7). One can directly check that the conditions in (4.5) are satisfied (as expected) and that the radius of the AdS_2 factor can now also be written

$$L^{-1} = \frac{1}{\sqrt{2}} \sum_i \bar{X}_i. \quad (4.15)$$

At this point we have shown that any solutions to the flow equations (4.12) preserve 1/16 of the supersymmetries, i.e. two Poincaré supersymmetries, which is enhanced to 1/8 supersymmetry for the AdS_2 fixed points. For there to be supersymmetry enhancement, one needs to have solutions to another of the conditions in (4.8), but this is not compatible with (4.13), (4.14).

In the three-dimensional moduli space of solutions, labelled by (f_1, f_2, f_3) , we have a two dimensional locus of supersymmetric solutions fixed by (4.14), which we have plotted in figure 4. Let us discuss a few special cases. Firstly, there are supersymmetric solutions when one of the f_a is set to zero. However, there are no supersymmetric solutions when two of the f_a are set to zero. In particular, there are no supersymmetric solutions with $U(1)^2 \times SU(2)^2$ symmetry that have e.g. $X_1 = X_2$, $X_3 = X_4$, $q^1 = q^2$ and $q^3 = q^4$, as noted in [18]. The AdS_2 solutions of minimal gauged supergravity with all f_a zero are not supersymmetric, as is well known.

Secondly, there are supersymmetric solutions when we set two of the f_a equal. Furthermore, there is a single supersymmetric solution when we set all of them to be equal, $f_1 = f_2 = f_3$. Specifically, in the $SU(3) \times U(1)^2$ invariant class solutions given in (4.6) we should take $q^1 = -3q^2$ (a condition that was also noted in [18]) with

$$\bar{X} = \left(-1 + \frac{2}{\sqrt{3}} \right)^{1/4}, \quad q^2 = \frac{1}{3\alpha} \sqrt{9 + 6\sqrt{3}}, \quad L^{-1} = 2 \left(9 + 6\sqrt{3} \right)^{1/4}. \quad (4.16)$$




Figure 4: The moduli space of supersymmetric magnetic $AdS_2 \times \mathbb{R}^2$ solutions in the f_1, f_2, f_3 space.

We show in appendix C that the uplifted $D = 11$ metric for this solution, using the formulae in [41], can be recast in the formalism of [51]. This provides a direct and very satisfying check on the supersymmetry of the solution.

4.3 Supersymmetric AdS_4 to $AdS_2 \times \mathbb{R}^2$ domain wall

We will be interested in constructing a supersymmetric domain wall solution that describes a flow from the AdS_4 vacuum to the fixed point (4.16) which preserves $U(1)^2 \times SU(3)$ in eleven dimensions. To construct the flow we truncate to the $SU(3)$ invariant sector by setting $\phi_a = \phi$ in (4.12) to obtain the first order system of equations

$$\begin{aligned} W' - \frac{1}{2\sqrt{2}}e^{-2U-\frac{3}{2}\phi} \left(e^{2U}(1+3e^{2\phi}) + \sqrt{9+6\sqrt{3}}(e^\phi - e^{3\phi}) \right) &= 0, \\ U' - \frac{1}{2\sqrt{2}}e^{-2U-\frac{3}{2}\phi} \left(e^{2U}(1+3e^{2\phi}) - \sqrt{9+6\sqrt{3}}(e^\phi - e^{3\phi}) \right) &= 0, \\ \phi' + \frac{1}{3\sqrt{2}}e^{-2U-\frac{3}{2}\phi} \left(-3e^{2U}(1-e^{2\phi}) + \sqrt{9+6\sqrt{3}}(e^\phi + 3e^{3\phi}) \right) &= 0. \end{aligned} \quad (4.17)$$

The expansion close to the $AdS_2 \times \mathbb{R}^2$ fixed point in the far IR is

$$\begin{aligned} W &= w_0 + L^{-1}\rho - 2\frac{3+\sqrt{3}}{3+2\sqrt{3}}c_{IR}e^{L^{-1}\rho} + \dots, \\ U &= c_{IR}e^{L^{-1}\rho} + \dots, \\ \phi &= -\frac{1}{4}\ln\left[3\left(7+4\sqrt{3}\right)\right] + \frac{2}{2+\sqrt{3}}c_{IR}e^{L^{-1}\rho} + \dots, \end{aligned} \quad (4.18)$$




Figure 5: We have plotted U' (blue) W' (purple) and ϕ (green) as functions of ρ for the $U(1)^2 \times SU(3)$ invariant supersymmetric domain wall solution interpolating between AdS_4 and $AdS_2 \times \mathbb{R}^2$.

with c_{IR} and w_0 being constants of integration. Setting the magnetic charges $q^i = 0$ in the flow equations (4.12), we recover the AdS_4 solution

$$W = U = R^{-1} \rho, \quad \phi = 0, \quad R^{-1} = \sqrt{2}. \quad (4.19)$$

Turning on a non-zero q^i triggers the following asymptotic expansion to the equations (4.17) given by

$$\begin{aligned} W &= R^{-1} \rho - \frac{3}{16} c_{UV}^2 e^{-2R^{-1} \rho} + \dots, \\ U &= R^{-1} \rho - \frac{3}{16} c_{UV}^2 e^{-2R^{-1} \rho} + \dots, \\ \phi &= c_{UV} e^{-R^{-1} \rho} + \left(2\sqrt{1 + \frac{2}{\sqrt{3}}} - \frac{c_{UV}^2}{2} \right) e^{-2R^{-1} \rho}, \end{aligned} \quad (4.20)$$

where c_{UV} is a constant of integration. This expansion corresponds to the operator dual to ϕ having a deformation as well as a VEV, and we see that both the deformation and the VEV are fixed by c_{UV} .

Using a shooting method we find that there is a solution to (4.12) with boundary conditions (4.18) and (4.20) with

$$w_0 = -0.47\dots, \quad c_{IR} = 0.26\dots, \quad c_{UV} = -1.71\dots, \quad (4.21)$$

as we have indicated in figure 5. This is the supersymmetric domain wall solution. This solution can be uplifted on S^7 , or an orbifold thereof, to $D = 11$ supergravity using the formulae in [41]. The uplifted solutions then describe the dual $d = 3$ SCFTs deformed by the presence of the magnetic field and also by the operators dual to ϕ . In particular, the supersymmetric $AdS_2 \times \mathbb{R}^2$ solutions describes the IR ground state at zero temperature.

5 Instabilities of magnetic $AdS_2 \times \mathbb{R}^2$ solutions

The instabilities for the magnetic $AdS_2 \times \mathbb{R}^2$ solutions that we constructed in section 4 are very similar to those that we have discussed for the $AdS_3 \times \mathbb{R}^2$ solutions in section 3. In this section, we will just present some illustrative calculations.

5.1 Spatially modulated instabilities of neutral scalars

We first investigate the possibility of spatially modulated instabilities of the neutral scalars ϕ_a for the magnetic $AdS_2 \times \mathbb{R}^2$ solutions given in (4.5). For simplicity we just analyse the one-dimensional subspace of $SU(3) \times U(1)^2$ solutions given in (4.6). In particular, we focus on the perturbation with

$$\begin{aligned} \delta\phi_1 &= 0, & \delta\phi_2 &= -\delta\phi_3 = \phi(t, \rho) \cos(kx_1), \\ \delta A^1 &= \delta A^2 = 0, & \delta A^3 &= -\delta A^4 = a(t, \rho) \sin(kx_1) dx_2. \end{aligned} \quad (5.1)$$

Defining the vector $\mathbf{v} = (\phi, a)$, the equations of motion for the Lagrangian (4.1) imply, at linear order,

$$(\square_{AdS_2} - L^2 M^2) \mathbf{v} = 0, \quad (5.2)$$

where the Laplacian is with respect to a unit radius AdS_2 and the mass matrix is

$$M^2 = \begin{pmatrix} k^2 + 2X^2 + 6X^{-2} & 8q^2 X^{-2} k \\ q^2 k & k^2 \end{pmatrix}. \quad (5.3)$$

The matrix M^2 has eigenvalues

$$m_{\pm}^2 = \frac{1}{X^2} \left[3 + k^2 \bar{X}^2 + \bar{X}^4 \pm \sqrt{8k^2 \bar{X}^2 (2 + \bar{X}^4) + (3 + \bar{X}^4)^2} \right]. \quad (5.4)$$

The branch m_-^2 develops a minimum at

$$k_{min}^2 = \frac{1}{8\bar{X}^2} \frac{55 + 58\bar{X}^4 + 15\bar{X}^8}{2 + \bar{X}^4}, \quad (5.5)$$

with the corresponding minimum satisfying

$$L^2 m_{min}^2 = -\frac{1}{48} \frac{(5 + 3\bar{X}^4)^2}{2 + 3\bar{X}^4 + \bar{X}^8}. \quad (5.6)$$

For $\bar{X} < \left(-1 + \frac{2}{\sqrt{3}}\right)^{1/4}$ the mass minimum violates the AdS_2 BF bound of $-1/4$ making the solution unstable.

It is worth noting that for the supersymmetric solution with $\bar{X} = \left(-1 + \frac{2}{\sqrt{3}}\right)^{1/4}$, the static mode given by

$$\phi(\rho) = c_1 e^{-\frac{\rho}{2L}} \cos(|k_{min}| x_1), \quad a(r) = c_2 e^{-\frac{\rho}{2L}} \sin(|k_{min}| x_1), \quad (5.7)$$

with $c_1^2/c_2^2 = 8\sqrt{-9 + 6\sqrt{3}}$, which saturates the BF bound, preserves the supersymmetries of the background. The sign choice of c_1 depends on the choice of α in the projector (4.11).

5.2 Instabilities of charged scalars

Recall that $SO(8)$ gauged supergravity has 70 scalar fields, parametrising the coset $E_{7(7)}/SU(8)$ and transforming as two **35** irreps of $SO(8)$. The scalars in one of these **35** irreps can be described by a unimodular 8×8 matrix T . The three neutral scalars we have been considering lie in this irrep. We next investigate possible instabilities of four charged fields lying in this irrep using the consistent truncation of $SO(8)$ gauge supergravity discussed in [45]. We should recall that $A^{there} = 2\sqrt{2}A^{here}$ and set their $g = 1/\sqrt{2}$. We then follow the earlier analysis in section 3.2 and for the scalar φ_1 (say) we find Landau levels with corresponding AdS_2 mass given by

$$m^2 = -2L^2 (X_1 X_2 + X_1 X_3 + X_1 X_4 - X_1^2 - (2n+1)|q_1|). \quad (5.8)$$

These modes violate the BF bound for a large parameter space of solutions. Indeed, for the lowest level, $n = 0$, the unstable regions are, roughly, the obvious generalisation of figure 3(b) to figure 4. In particular, these modes now intersect the locus of supersymmetric solutions in a one-dimensional sub-locus.

Finally, we note that there will be additional instabilities for the other scalars in the 8×8 matrix T and these will mix with the gauge fields and the analysis will mirror the analysis that we carried out in section (3.3).

6 Electric Solutions

In this section we construct new electric $AdS_2 \times \mathbb{R}^2$ and $AdS_2 \times \mathbb{R}^3$ solutions.

6.1 Electric $AdS_2 \times \mathbb{R}^2$ solutions

The equations of motion of the $U(1)^4$ truncation of $D = 4$ $SO(8)$ gauged supergravity (4.1) are invariant under the electric-magnetic duality transformation

$$F^i \rightarrow X_i^{-2} * F^i, \quad \phi_a \rightarrow -\phi_a, \quad (6.1)$$

with the metric unchanged. We can use this symmetry to immediately obtain electric analogues of the magnetic solutions that we presented in section 4.

Starting with (4.3), (4.5) we obtain electric $AdS_2 \times \mathbb{R}^2$ solutions, which we can write as

$$\begin{aligned} ds^2 &= L^2 ds^2(AdS_2) + dx_1^2 + dx_2^2, \\ F^i &= \frac{1}{2} Q^i L^2 Vol(AdS_2), \\ \phi_1 &= f_1, \quad \phi_2 = f_2, \quad \phi_3 = f_3, \end{aligned} \tag{6.2}$$

where

$$(Q^i)^2 = \bar{X}_i^3 \sum_{j \neq i} \bar{X}_j, \quad L^{-2} = -2V(\bar{X}_i), \tag{6.3}$$

and, as before,

$$\bar{X}_1 = e^{\frac{1}{2}(-f_1-f_2-f_3)}, \quad \bar{X}_2 = e^{\frac{1}{2}(-f_1+f_2+f_3)}, \quad \bar{X}_3 = e^{\frac{1}{2}(f_1-f_2+f_3)}, \quad \bar{X}_4 = e^{\frac{1}{2}(f_1+f_2-f_3)}. \tag{6.4}$$

None of the solutions preserve the supersymmetry transformations given in (4.10). For the special case of the electric $AdS_2 \times \mathbb{R}^2$ solution of minimal gauged supergravity see [40] for a discussion on instabilities. An analysis for other solutions will be carried out elsewhere.

Starting with the supersymmetric domain wall solution that we presented in section 4.3, we can use the duality transformation (6.1) to immediately obtain an electric domain wall solution that interpolates between AdS_4 in the UV and an electric $AdS_2 \times \mathbb{R}^2$ solution in the IR. Note that despite the domain wall not preserving supersymmetry it solves first order flow equations.

6.2 Electric $AdS_2 \times \mathbb{R}^3$ solutions

We now consider electric $AdS_2 \times \mathbb{R}^3$ solutions of $D = 5$ $SO(6)$ gauged supergravity. By direct construction we find

$$\begin{aligned} ds^2 &= L^2 ds^2(AdS_2) + dx_1^2 + dx_2^2 + dx_3^2, \\ F^i &= 2Q^i L^2 Vol(AdS_2), \quad \phi_1 = f_1, \quad \phi_2 = f_2, \end{aligned} \tag{6.5}$$

where f_a are constants,

$$(Q^i)^2 = \bar{X}_i^3 \sum_{j \neq i} \bar{X}_j, \quad L^{-2} = -V(\bar{X}_i), \tag{6.6}$$

and $\bar{X}_i$ are the on-shell values

$$\bar{X}_1 = e^{-\frac{1}{\sqrt{6}}f_1 - \frac{1}{\sqrt{2}}f_2}, \quad \bar{X}_2 = e^{-\frac{1}{\sqrt{6}}f_1 + \frac{1}{\sqrt{2}}f_2}, \quad \bar{X}_3 = e^{\frac{2}{\sqrt{6}}f_1}. \quad (6.7)$$

When $f_2 = 0$, for example, these are solutions to Romans' theory, and actually were already presented in [6] and further discussed in appendix B of [22]. When $f_1 = f_2 = 0$ we obtain the standard $AdS_2 \times \mathbb{R}^3$ solution of minimal gauged supergravity which is the near horizon limit of the usual electrically charged AdS-RN black brane solution. Note that the solutions do not preserve the supersymmetry (2.9); within Romans' theory this was shown in [6].

It was shown in [22] that the electrically charged $AdS_2 \times \mathbb{R}^3$ solutions in Romans' theory all suffer from instabilities corresponding to helical p -wave superconductors. A more detailed stability analysis of all solutions will be carried out elsewhere.

Acknowledgements

We would like to thank Nikolay Bobev and Joe Polchinski for helpful discussions. AD is supported by an EPSRC Postdoctoral Fellowship. JPG is supported by a Royal Society Wolfson Award. CP is supported by an I.K.Y. Scholarship.

A Charged mode analysis for the truncation [43]

A consistent truncation that supplements the $U(1)^3 \subset SO(6)$ truncation (2.2) with four complex scalars ζ_m , $m = 1, 2, 3, 4$ was given in [43]. After writing $\zeta_m = \tanh(\gamma_m)e^{i\theta_m}$, the Lagrangian is given in eq. (2.7) of [43] and, to make contact with our notation, one should set $\varphi_m^{there} = \gamma_m^{here}$, $g^{there} = 2$ and also identify the gauge fields via $A^{i(there)} = A^{i(here)}/2$. It is straightforward to see that after expanding the equations of motion around the $AdS_3 \times \mathbb{R}^2$ solutions (2.4), the fields θ_m are all massless. The analysis for the charged modes γ_m is very similar to that in section 3.2. For example, for $m = 1$ we find that the lowest mass mode is obtained by writing

$$\gamma_1 = e^{-\frac{|q_1+q_2-q_3|}{4}(x_1^2+x_2^2)} \sigma_1(t, r, z), \quad (A.1)$$

giving

$$(\square_{AdS_3} - L^2 m_{\sigma_1}^2) \sigma_1 = 0, \quad (A.2)$$

with

$$m_{\sigma_1}^2 = |q_1 + q_2 - q_3| + \sum_i \bar{X}_i^2 - 2 \sum_i \bar{X}_i^{-1}. \quad (A.3)$$

Over the moduli space of $AdS_3 \times \mathbb{R}^2$ slutions we find that the minimum value is $L^2 m_{\sigma_1}^2 \approx -0.704$ and does not violate the BF bound.

B The mixed charged modes

Here we provide some details of the calculations we carried out for section 3.3. We consider the perturbation t, a about the background $AdS_3 \times \mathbb{R}^2$ solution (2.4),(2.5) defined by

$$T = \tilde{T} + t, \quad A = \tilde{A} + a, \quad (B.1)$$

with

$$\tilde{T}_{I\bar{I}} = X_I, \quad \tilde{A}_{I\bar{I}} = q_I (\bar{z} dz - z d\bar{z}). \quad (B.2)$$

It is is useful to note that at leading order in the perturbation we have $T^{-1} = \tilde{T}^{-1} - \tilde{T}^{-1} t \tilde{T}^{-1}$. Furthermore, the linearised expression of the field strengths and some covariant derivatives are given by

$$\begin{aligned} \delta(F_{IJ}) &= da_{IJ} + \left(\tilde{A}_{I\bar{I}} + \tilde{A}_{J\bar{J}} \right) \wedge a_{IJ}, \\ \delta(F_{I\bar{J}}) &= da_{I\bar{J}} + \left(\tilde{A}_{I\bar{I}} - \tilde{A}_{J\bar{J}} \right) \wedge a_{I\bar{J}}, \\ \delta(DT_{IJ}) &= dt_{IJ} + \left(\tilde{A}_{I\bar{I}} + \tilde{A}_{J\bar{J}} \right) t_{IJ} + g (X_J - X_I) a_{IJ}, \\ \delta(DT_{I\bar{J}}) &= dt_{I\bar{J}} + \left(\tilde{A}_{I\bar{I}} - \tilde{A}_{J\bar{J}} \right) t_{I\bar{J}} + g (X_J - X_I) a_{I\bar{J}}, \\ \delta(D * F_{IJ}) &= d * \delta(F_{IJ}) + \left(\tilde{A}_{I\bar{I}} + \tilde{A}_{J\bar{J}} \right) \wedge * \delta(F_{IJ}) + * \left(\tilde{F}_{I\bar{I}} + \tilde{F}_{J\bar{J}} \right) \wedge a_{IJ}, \\ \delta(D * DT_{IJ}) &= d * \delta(Dt_{IJ}) + \left(\tilde{A}_{I\bar{I}} + \tilde{A}_{J\bar{J}} \right) \wedge * \delta(Dt_{IJ}). \end{aligned} \quad (B.3)$$

We will only provide details concerning the IJ components of the equations of motion (3.15), with $I \neq J$. The case of $I\bar{J}$ is very similar. At linearised order we have

$$\delta(D * F_{IJ}) + \left(X_I^{-1} * \tilde{F}_{I\bar{I}} - X_J^{-1} * \tilde{F}_{J\bar{J}} \right) \wedge \delta(Dt_{IJ}) = - (X_J - X_I) * \delta(Dt_{IJ}), \quad (B.4)$$

$$- \delta(D * Dt_{IJ}) = 4q_I^{-1} q_J^{-1} t_{IJ} + * \left(X_J^{-1} \tilde{F}_{J\bar{J}} - X_I^{-1} \tilde{F}_{I\bar{I}} \right) \wedge \delta(F_{IJ}). \quad (B.5)$$

For the gauge fields we take

$$a_{IJ} = a_{IJ}^1 dz + a_{IJ}^2 d\bar{z}, \quad (B.6)$$

and, after defining

$$\omega_{IJ} = q_I + q_J, \quad (\text{B.7})$$

we find

$$\begin{aligned} \delta(F_{IJ}) &= da_{IJ}^1 \wedge dz + da_{IJ}^2 \wedge d\bar{z} - (\partial_{\bar{z}} - z\omega_{IJ}) a_{IJ}^1 dz \wedge d\bar{z} + (\partial_z + \bar{z}\omega_{IJ}) a_{IJ}^2 dz \wedge d\bar{z}, \\ \delta(Dt_{IJ}) &= dt_{IJ} + (\partial_z + \bar{z}\omega_{IJ}) t_{IJ} dz + (\partial_{\bar{z}} - z\omega_{IJ}) t_{IJ} d\bar{z} + (X_J - X_I) (a_{IJ}^1 dz + a_{IJ}^2 d\bar{z}), \end{aligned} \quad (\text{B.8})$$

where d is the exterior derivative on AdS_3 . We can obtain analogous expressions for $\delta(D * F_{IJ})$ and $\delta(D * Dt_{IJ})$ using B.3, which we then substitute into the equations of motion (B.4) and (B.5).

From (B.4) we are led to impose the constraint

$$(\partial_{\bar{z}} - \omega_{IJ}z) a_{IJ}^1 + (\partial_z + \omega_{IJ}\bar{z}) a_{IJ}^2 = -(X_J - X_I) t_{IJ}. \quad (\text{B.9})$$

We also obtain

$$\begin{aligned} &-L^{-2}\square_{AdS_3} a_{IJ}^1 + 2(-\partial_z\partial_{\bar{z}} + \omega_{IJ}^2 z\bar{z} + 2\omega_{IJ} + \omega_{IJ}(z\partial_z - \bar{z}\partial_{\bar{z}})) a_{IJ}^1 \\ &+ 2(q_I^{-1} - q_J^{-1}) [(\partial_z + \omega_{IJ}\bar{z}) t_{IJ} + (X_J - X_I) a_{IJ}^1] = -(X_J - X_I)^2 a_{IJ}^1, \end{aligned} \quad (\text{B.10})$$

and

$$\begin{aligned} &L^{-2}\square_{AdS_3} a_{IJ}^2 + 2(\partial_z\partial_{\bar{z}} - \omega_{IJ}^2 z\bar{z} + 2\omega_{IJ} - \omega_{IJ}(z\partial_z - \bar{z}\partial_{\bar{z}})) a_{IJ}^2 \\ &+ 2(q_I^{-1} - q_J^{-1}) [(\partial_{\bar{z}} - \omega_{IJ}z) t_{IJ} + (X_J - X_I) a_{IJ}^2] = (X_J - X_I)^2 a_{IJ}^2, \end{aligned} \quad (\text{B.11})$$

where $\square_{AdS_3}$ is the Laplacian on a unit radius AdS_3 . Similarly, from (B.5) we obtain

$$\begin{aligned} &L^{-2}\square_{AdS_3} t_{IJ} + 2(\partial_z\partial_{\bar{z}} - \omega_{IJ}^2 z\bar{z} + \omega_{IJ}(\bar{z}\partial_{\bar{z}} - z\partial_z)) t_{IJ} - (X_J - X_I)^2 t_{IJ} = \\ &-4q_I^{-1}q_J^{-1}t_{IJ} + 2(q_I^{-1} - q_J^{-1}) [(-\partial_{\bar{z}} + \omega_{IJ}z) a_{IJ}^1 + (\partial_z + \omega_{IJ}\bar{z}) a_{IJ}^2]. \end{aligned} \quad (\text{B.12})$$

We now observe that because of the constraint (B.9), the three equations (B.10)-(B.12) are not independent. Indeed acting on equation (B.10) by $(-\partial_{\bar{z}} + \omega_{IJ}z)$, on equation (B.11) by $(\partial_z + \omega_{IJ}\bar{z})$ and adding one can show that equation (B.12) is satisfied.

To continue with the analysis, we need to fix the sign of $\omega \equiv \omega_{IJ}$. We first take $\omega > 0$. For this case we can keep equation (B.11) and (B.12) which we write as

$$\begin{aligned} &L^{-2}\square_{AdS_3} t_{IJ} + 2(\partial_z\partial_{\bar{z}} - \omega^2 z\bar{z} + \omega(\bar{z}\partial_{\bar{z}} - z\partial_z)) t_{IJ} - (X_J - X_I)^2 t_{IJ} = \\ &-4q_I^{-1}q_J^{-1}t_{IJ} + 2(q_I^{-1} - q_J^{-1}) [(X_J - X_I) t_{IJ} + 2(\partial_z + \omega\bar{z}) a_{IJ}^2]. \end{aligned} \quad (\text{B.13})$$

Next we introduce the ladder operators

$$\begin{aligned} a &= \frac{1}{\sqrt{2\omega}} (\partial_{\bar{z}} + \omega z), & a^\dagger &= \frac{1}{\sqrt{2\omega}} (-\partial_z + \omega \bar{z}), \\ b &= \frac{1}{\sqrt{2\omega}} (\partial_z + \omega \bar{z}), & b^\dagger &= \frac{1}{\sqrt{2\omega}} (-\partial_{\bar{z}} + \omega z), \end{aligned} \quad (\text{B.14})$$

which can be checked to satisfy the algebra

$$[a, a^\dagger] = 1, \quad [b, b^\dagger] = 1, \quad (\text{B.15})$$

and the rest of the commutators being trivial. Note that we have

$$\begin{aligned} -\partial_z \partial_{\bar{z}} + \omega^2 \bar{z} z &= \omega (a^\dagger a + b^\dagger b + 1), \\ \bar{z} \partial_{\bar{z}} - z \partial_z &= a^\dagger a - b^\dagger b. \end{aligned} \quad (\text{B.16})$$

In terms of these operators, equations (B.11) and (B.13) take the form

$$\begin{aligned} L^{-2} \square_{AdS_3} a_{IJ}^2 - 2\omega (2b^\dagger b - 1) a_{IJ}^2 \\ + 2 (q_I^{-1} - q_J^{-1}) \left[-\sqrt{2\omega} b^\dagger t_{IJ} + (X_J - X_I) a_{IJ}^2 \right] - (X_J - X_I)^2 a_{IJ}^2 = 0 \end{aligned} \quad (\text{B.17})$$

and

$$\begin{aligned} L^{-2} \square_{AdS_3} t_{IJ} - 2\omega (2b^\dagger b + 1) t_{IJ} - (X_J - X_I)^2 t_{IJ} + 4q_I^{-1} q_J^{-1} t_{IJ} \\ - 2 (q_I^{-1} - q_J^{-1}) \left[(X_J - X_I) t_{IJ} + 2\sqrt{2\omega} b a_{IJ}^2 \right] = 0. \end{aligned} \quad (\text{B.18})$$

To reduce the problem to modes on the AdS_3 space we introduce the ground state

$$L_{0,0} = \left(\frac{\omega}{\pi} \right)^{1/2} e^{-\omega z \bar{z}} \quad (\text{B.19})$$

and the complete set of functions

$$L_{n,m}(z, \bar{z}) = \frac{(b^\dagger)^n (a^\dagger)^m}{\sqrt{n!} \sqrt{m!}} L_{0,0}(z, \bar{z}), \quad m, n > 0. \quad (\text{B.20})$$

We use these to write the expansions

$$t_{IJ} = \sum_{n,m} f_{IJ}^{n,m} L_{n,m}(z, \bar{z}), \quad a_{IJ}^2 = \sum_{n,m} g_{IJ}^{n,m} L_{n,m}(z, \bar{z}), \quad (\text{B.21})$$

with f and g defined on AdS_3 . From equations (B.17) and (B.18) we see that the modes $g_{IJ}^{0,m}$ decouple and they have an AdS_3 mass

$$m_{0,m}^2 = -2\omega - 2 (q_I^{-1} - q_J^{-1}) (X_J - X_I) + (X_J - X_I)^2. \quad (\text{B.22})$$

For the rest of the modes, we see that $g_{IJ}^{n+1,m}$ mix with $f_{IJ}^{n,m}$ for $n \geq 0$ with mass matrix

$$M^2 = \begin{pmatrix} 2\omega(2n+1) - 4q_I^{-1}q_J^{-1} + 2W_{IJ}V_{JI} + V_{IJ}^2 & 4\sqrt{2\omega}W_{IJ}(n+1)^{1/2} \\ 2\sqrt{2\omega}W_{IJ}(n+1)^{1/2} & 2\omega(2n+1) - 2W_{IJ}V_{JI} + V_{IJ}^2 \end{pmatrix}, \quad (\text{B.23})$$

where we set

$$W_{IJ} = q_I^{-1} - q_J^{-1}, \quad V_{IJ} = X_I - X_J. \quad (\text{B.24})$$

Note that our results differ from those presented in the first two lines of eq. (6.7) and eq. (6.12) of [18] because the mixing of the charged scalars and vectors was not taken into account in that reference¹¹.

When $\omega < 0$, in order to get the zero modes, we should keep equations (B.10) and (B.12). For the ladder operators we should take $\sqrt{2\omega} \rightarrow \sqrt{2|\omega|}$ and $\omega \rightarrow -\omega$ in (B.14). A very similar analysis then ensues and we obtain (B.22) and (B.23) after substituting $\omega \rightarrow |\omega|$ and also $W_{IJ} \rightarrow -W_{IJ}$.

Finally, let us consider $\omega = 0$. This occurs along the three lines in figure 1 with $X_I = X_J$. The equations (B.9) - (B.12) then simplify and we are essentially led back to the mass matrix that we saw for spatially modulated neutral scalars in (3.3).

C Construction of SUSY $AdS_2 \times \mathbb{R}^2$ solutions

Recall [51] that supersymmetric AdS_2 solutions of $D = 11$ supergravity with purely electric four-form flux, generically dual to CFTs with two (Poincaré) supersymmetries, can be obtained from an eight dimensional Kähler metric, ds_8^2 , whose Ricci tensor satisfies

$$\square_8 R - \frac{1}{2}R^2 + R_{ij}R^{ij} = 0. \quad (\text{C.1})$$

The $D = 11$ metric has the form

$$ds^2 = e^{2A} [ds^2(AdS_2) + e^{-3A} ds_8^2 + (dz + P)^2], \quad (\text{C.2})$$

where $dP = \mathcal{R}$, where $\mathcal{R}$ is the Ricci-form, and $e^{-3A} = \frac{1}{2}R$.

¹¹Our zero modes (B.22) only involve the gauge fields and the fact that our results differ from eq. (6.12) of [18] after setting their $n = 0$, which also just involve the gauge-fields, can be traced back to the fact that our equation of motion (B.4) for $t_{IJ} = 0$ does not come from the Lagrangian given in eq. (6.11) of [18].

Following [5] we start with an ansatz for an eight dimensional Kähler metric given by

$$ds_8^2 = \frac{dy^2}{U} + y^2 U (D\phi + A)^2 + y^2 ds^2(\mathbb{CP}^2) + (ay^2 + b) ds^2(\mathbb{R}^2), \quad (\text{C.3})$$

with $dD\phi = 2 J_{\mathbb{CP}^2}$, where $J_{\mathbb{CP}^2}$ is the Kähler form on $\mathbb{CP}^2$, and $dA = 2a J_{\mathbb{R}^2}$, where $J_{\mathbb{R}^2} = \text{Vol}(\mathbb{R}^2)$ and a is a constant, and $U = U(y)$. For the corresponding holomorphic 2-form and 1-form on $\mathbb{CP}^2$ and $\mathbb{R}^2$ we have

$$\begin{aligned} d\Omega_{\mathbb{CP}^2} &= i P_{\mathbb{CP}^2} \wedge \Omega_{\mathbb{CP}^2}, \quad dP_{\mathbb{CP}^2} = 2l J_{\mathbb{CP}^2}, \\ d\Omega_{\mathbb{R}^2} &= 0, \end{aligned} \quad (\text{C.4})$$

where l is another (positive) constant. The Kähler-form and holomorphic 4-form for the eight dimensional space can now be written

$$\begin{aligned} J &= y dy \wedge (D\phi + A) + y^2 J_{\mathbb{CP}^2} + (ay^2 + b) J_{\mathbb{R}^2}, \\ \Omega_4 &= e^{i\phi} y^2 \sqrt{ay^2 + b} \left[\frac{dy}{\sqrt{U}} + iy\sqrt{U} (D\phi + A) \right] \wedge \Omega_{\mathbb{CP}^2} \wedge \Omega_{\mathbb{R}^2}. \end{aligned} \quad (\text{C.5})$$

We can easily show that

$$d\Omega_4 = i P \wedge \Omega_4, \quad P = l D\phi - g (D\phi + A), \quad (\text{C.6})$$

where

$$g = 3U + \frac{ay^2 U}{ay^2 + b} + \frac{yU'}{2}. \quad (\text{C.7})$$

The Ricci form for the eight dimensional space is given by

$$\mathcal{R} = dP = 2(l - g) J_{\mathbb{CP}^2} - 2ag J_{\mathbb{R}^2} - g' dy \wedge (D\phi + A). \quad (\text{C.8})$$

In order to get an $AdS_2 \times \mathbb{R}^2$ factor in (C.2) we now require that the Ricci scalar of the eight dimensional space satisfies

$$R = \frac{W}{ay^2 + b}, \quad (\text{C.9})$$

for some constant W . The resulting second order equation for U gives the solution

$$U = \frac{1}{48} \frac{1}{ay^2 + b} \left(16bl + 8aly^2 - Wy^2 + \frac{c_1}{y^6} + \frac{c_2}{y^4} \right), \quad (\text{C.10})$$

where c_i are two constants of integration. One can check that in order to solve (C.1) we need to set $c_1 = c_2 = 0$ and we then find two solutions

$$W = -4 \left(1 \mp \sqrt{3} \right) al \quad \Rightarrow \quad U = \frac{l}{12} \frac{4b + (3 \mp \sqrt{3}) ay^2}{b + ay^2}. \quad (\text{C.11})$$

Let us now continue with the solution with the lower sign. We take $a < 0$ and change coordinates via $y = \frac{2}{\sqrt{3+\sqrt{3}}} \sqrt{-\frac{b}{a}} \sin \xi$ and we will take $0 < \xi < \pi/2$. We find

$$\begin{aligned} U &= l \left(1 + \frac{1}{\sqrt{3}} \right) \frac{\cos^2 \xi}{1 + \sqrt{3} + 2 \cos 2\xi}, \\ R_8 &= -8 \frac{al (3 + 2\sqrt{3})}{b} \frac{1}{1 + \sqrt{3} + 2 \cos 2\xi}, \\ g &= l \frac{3 + \sqrt{3}}{3} \frac{1 + 2 \cos 2\xi}{1 + \sqrt{3} + 2 \cos 2\xi}. \end{aligned} \quad (\text{C.12})$$

We can now assemble the $D = 11$ metric using (C.2) and, after setting $l = 3$, find

$$\begin{aligned} ds^2 &= e^{2A} L^{-2} \left\{ L^2 ds^2(AdS_2) + \frac{L^2 W}{2} ds^2(\mathbb{R}^2) \right. \\ &\quad \left. + 2\bar{X}^2 \left(d\xi^2 + \frac{\sin^2 \xi}{\bar{X}^3 \Delta} [ds^2(CP^2) + (D\tilde{\phi} + \frac{1 + \sqrt{3}}{4} A)^2] + \frac{\bar{X} \cos^2 \xi}{\Delta} \frac{1}{8\sqrt{3}\bar{X}^4} (dz - 3A)^2 \right) \right\} \end{aligned} \quad (\text{C.13})$$

where $\bar{X}$ is as in (4.16) and

$$\Delta = \frac{1}{\bar{X}^3} (\cos^2 \xi + \bar{X}^4 \sin^2 \xi). \quad (\text{C.14})$$

One can now check that if we set

$$-\frac{a}{b} = \frac{4}{3^{1/4}(1 + \sqrt{3})} \quad (\text{C.15})$$

and rescale $(L^2 W/2) ds^2(\mathbb{R}^2) \rightarrow ds^2(\mathbb{R}^2)$ then we precisely obtain the uplift of the solution (4.16) using the formulae in [41] (setting $g^2 = 1/2$ in eq. (3.8) of [41] and identifying $(F^i)^{there} = 2\sqrt{2}(F^i)^{here}$).

References

- [1] S. A. Hartnoll and P. Kovtun, “Hall conductivity from dyonic black holes,” *Phys. Rev.* **D76** (2007) 066001, [arXiv:0704.1160 \[hep-th\]](#).
- [2] S. A. Hartnoll, P. K. Kovtun, M. Muller, and S. Sachdev, “Theory of the Nernst effect near quantum phase transitions in condensed matter, and in dyonic black holes,” *Phys. Rev.* **B76** (2007) 144502, [arXiv:0706.3215 \[cond-mat.str-el\]](#).

- [3] S. A. Hartnoll and C. P. Herzog, “Ohm’s Law at strong coupling: S duality and the cyclotron resonance,” *Phys. Rev.* **D76** (2007) 106012, [arXiv:0706.3228 \[hep-th\]](#).
- [4] T. Albash and C. V. Johnson, “A Holographic Superconductor in an External Magnetic Field,” *JHEP* **0809** (2008) 121, [arXiv:0804.3466 \[hep-th\]](#).
- [5] J. P. Gauntlett and O. Varela, “D=5 $SU(2) \times U(1)$ Gauged Supergravity from D=11 Supergravity,” *JHEP* **02** (2008) 083, [arXiv:0712.3560 \[hep-th\]](#).
- [6] L. J. Romans, “Gauged N=4 supergravities in five-dimensions and their magnetovac backgrounds,” *Nucl. Phys.* **B267** (1986) 433.
- [7] H. Lin, O. Lunin, and J. M. Maldacena, “Bubbling AdS space and 1/2 BPS geometries,” *JHEP* **10** (2004) 025, [arXiv:hep-th/0409174](#).
- [8] D. Gaiotto and J. Maldacena, “The gravity duals of N=2 superconformal field theories,” [arXiv:0904.4466 \[hep-th\]](#).
- [9] E. D’Hoker and P. Kraus, “Charged Magnetic Brane Solutions in AdS_5 and the fate of the third law of thermodynamics,” *JHEP* **03** (2010) 095, [arXiv:0911.4518 \[hep-th\]](#).
- [10] E. D’Hoker and P. Kraus, “Holographic Metamagnetism, Quantum Criticality, and Crossover Behavior,” *JHEP* **05** (2010) 083, [arXiv:1003.1302 \[hep-th\]](#).
- [11] E. D’Hoker and P. Kraus, “Magnetic Field Induced Quantum Criticality via new Asymptotically AdS_5 Solutions,” *Class. Quant. Grav.* **27** (2010) 215022, [arXiv:1006.2573 \[hep-th\]](#).
- [12] E. D’Hoker and P. Kraus, “Magnetic Brane Solutions in AdS,” *JHEP* **10** (2009) 088, [arXiv:0908.3875 \[hep-th\]](#).
- [13] J. P. Gauntlett, D. Martelli, J. Sparks, and D. Waldram, “Supersymmetric AdS(5) solutions of type IIB supergravity,” *Class. Quant. Grav.* **23** (2006) 4693–4718, [arXiv:hep-th/0510125](#).
- [14] J. P. Gauntlett, D. Martelli, J. Sparks, and D. Waldram, “Supersymmetric AdS(5) solutions of M-theory,” *Class. Quant. Grav.* **21** (2004) 4335–4366, [arXiv:hep-th/0402153](#).

- [15] J. P. Gauntlett, E. O. Colgain, and O. Varela, “Properties of some conformal field theories with M-theory duals,” *JHEP* **0702** (2007) 049, [arXiv:hep-th/0611219](#) [[hep-th](#)].
- [16] J. P. Gauntlett and O. Varela, “Consistent Kaluza-Klein Reductions for General Supersymmetric AdS Solutions,” *Phys. Rev.* **D76** (2007) 126007, [arXiv:0707.2315](#) [[hep-th](#)].
- [17] A. Almuhairi, “AdS₃ and AdS₂ Magnetic Brane Solutions,” [arXiv:1011.1266](#) [[hep-th](#)].
- [18] A. Almuhairi and J. Polchinski, “Magnetic $AdS \times R^2$: Supersymmetry and stability,” [arXiv:1108.1213](#) [[hep-th](#)].
- [19] S. Nakamura, H. Ooguri, and C.-S. Park, “Gravity Dual of Spatially Modulated Phase,” *Phys. Rev.* **D81** (2010) 044018, [arXiv:0911.0679](#) [[hep-th](#)].
- [20] A. Donos and J. P. Gauntlett, “Holographic striped phases,” *JHEP* **08** (2011) 140, [arXiv:1106.2004](#) [[hep-th](#)].
- [21] O. Bergman, N. Jokela, G. Lifschytz, and M. Lippert, “Striped instability of a holographic Fermi-like liquid,” [arXiv:1106.3883](#) [[hep-th](#)].
- [22] A. Donos and J. P. Gauntlett, “Holographic helical superconductors,” [arXiv:1109.3866](#) [[hep-th](#)].
- [23] M. Ammon, J. Erdmenger, P. Kerner, and M. Strydom, “Black Hole Instability Induced by a Magnetic Field,” [arXiv:1106.4551](#) [[hep-th](#)].
- [24] M. N. Chernodub, “Superconductivity of QCD vacuum in strong magnetic field,” *Phys. Rev.* **D82** (2010) 085011, [arXiv:1008.1055](#) [[hep-ph](#)].
- [25] N. K. Nielsen and P. Olesen, “An Unstable Yang-Mills Field Mode,” *Nucl. Phys.* **B144** (1978) 376.
- [26] J. Ambjorn and P. Olesen, “On the Formation of a Random Color Magnetic Quantum Liquid in QCD,” *Nucl. Phys.* **B170** (1980) 60.
- [27] M. Rasolt and Z. Tesanovic, “Theoretical aspects of superconductivity in very high magnetic fields,” *Rev. Mod. Phys.* **64** (1992) 709–754.

- [28] D. Aoki, T. D. Matsuda, V. Taufour, E. Hassinger, G. Knebel, and J. Flouquet, “Superconductivity Reinforced by Magnetic Field and the Magnetic Instability in Uranium Ferromagnets,” *J. Phys. Soc. Jpn.* **78** (2009) 113709, [arXiv:1012.1987 \[cond-mat\]](#).
- [29] S.-S. Lee, “A Non-Fermi Liquid from a Charged Black Hole: A Critical Fermi Ball,” *Phys.Rev.* **D79** (2009) 086006, [arXiv:0809.3402 \[hep-th\]](#).
- [30] H. Liu, J. McGreevy, and D. Vegh, “Non-Fermi liquids from holography,” *Phys.Rev.* **D83** (2011) 065029, [arXiv:0903.2477 \[hep-th\]](#).
- [31] M. Cubrovic, J. Zaanen, and K. Schalm, “String Theory, Quantum Phase Transitions and the Emergent Fermi-Liquid,” *Science* **325** (2009) 439–444, [arXiv:0904.1993 \[hep-th\]](#).
- [32] T. Faulkner, H. Liu, J. McGreevy, and D. Vegh, “Emergent quantum criticality, Fermi surfaces, and $AdS(2)$,” *Phys.Rev.* **D83** (2011) 125002, [arXiv:0907.2694 \[hep-th\]](#).
- [33] T. Albash and C. V. Johnson, “Holographic Aspects of Fermi Liquids in a Background Magnetic Field,” *J. Phys.* **A43** (2010) 345405, [arXiv:0907.5406 \[hep-th\]](#).
- [34] P. Basu, J. He, A. Mukherjee, and H.-H. Shieh, “Holographic Non-Fermi Liquid in a Background Magnetic Field,” *Phys. Rev.* **D82** (2010) 044036, [arXiv:0908.1436 \[hep-th\]](#).
- [35] A. Donos, J. P. Gauntlett, and C. Pantelidou, “Spatially modulated instabilities of magnetic black branes,” [arXiv:1109.0471 \[hep-th\]](#).
- [36] S. L. Cacciatori and D. Klemm, “Supersymmetric AdS_4 black holes and attractors,” *JHEP* **01** (2010) 085, [arXiv:0911.4926 \[hep-th\]](#).
- [37] G. Dall’Agata and A. Gnechchi, “Flow equations and attractors for black holes in $N = 2$ $U(1)$ gauged supergravity,” *JHEP* **03** (2011) 037, [arXiv:1012.3756 \[hep-th\]](#).
- [38] S. Barisch, G. L. Cardoso, M. Haack, S. Nampuri, and N. A. Obers, “Nernst branes in gauged supergravity,” *JHEP* **11** (2011) 090, [arXiv:1108.0296 \[hep-th\]](#).

- [39] I. R. Klebanov, S. S. Pufu, and T. Tesileanu, “Membranes with Topological Charge and AdS₄/CFT₃ Correspondence,” *Phys. Rev.* **D81** (2010) 125011, [arXiv:1004.0413 \[hep-th\]](#).
- [40] F. Denef and S. A. Hartnoll, “Landscape of superconducting membranes,” *Phys. Rev.* **D79** (2009) 126008, [arXiv:0901.1160 \[hep-th\]](#).
- [41] M. Cvetič *et al.*, “Embedding AdS black holes in ten and eleven dimensions,” *Nucl. Phys.* **B558** (1999) 96–126, [arXiv:hep-th/9903214](#).
- [42] S. L. Cacciatori, D. Klemm, and W. A. Sabra, “Supersymmetric domain walls and strings in D = 5 gauged supergravity coupled to vector multiplets,” *JHEP* **03** (2003) 023, [arXiv:hep-th/0302218](#).
- [43] N. Bobev, A. Kundu, K. Pilch, and N. P. Warner, “Supersymmetric Charged Clouds in AdS₅,” *JHEP* **03** (2011) 070, [arXiv:1005.3552 \[hep-th\]](#).
- [44] M. Cvetič, H. Lu, C. Pope, A. Sadrzadeh, and T. A. Tran, “Consistent SO(6) reduction of type IIB supergravity on S¹ × S⁴,” *Nucl. Phys.* **B586** (2000) 275–286, [arXiv:hep-th/0003103 \[hep-th\]](#).
- [45] Z. W. Chong, H. Lu, and C. N. Pope, “BPS geometries and AdS bubbles,” *Phys. Lett.* **B614** (2005) 96–103, [arXiv:hep-th/0412221](#).
- [46] J. T. Liu, H. Lu, C. N. Pope, and J. F. Vazquez-Poritz, “New supersymmetric solutions of N=2, D=5 gauged supergravity with hyperscalars,” *JHEP* **10** (2007) 093, [arXiv:0705.2234 \[hep-th\]](#).
- [47] M. J. Duff and J. T. Liu, “Anti-de Sitter black holes in gauged N = 8 supergravity,” *Nucl. Phys.* **B554** (1999) 237–253, [arXiv:hep-th/9901149](#).
- [48] N. P. Warner, “Some new extrema of the scalar potential of gauged N=8 supergravity,” *Phys. Lett.* **B128** (1983) 169.
- [49] N. Bobev, N. Halmagyi, K. Pilch, and N. P. Warner, “Supergravity Instabilities of Non-Supersymmetric Quantum Critical Points,” *Class. Quant. Grav.* **27** (2010) 235013, [arXiv:1006.2546](#).
- [50] J. P. Gauntlett, O. A. P. Mac Conamhna, T. Mateos, and D. Waldram, “AdS spacetimes from wrapped M5 branes,” *JHEP* **11** (2006) 053, [arXiv:hep-th/0605146](#).

- [51] N. Kim and J.-D. Park, “Comments on AdS(2) solutions of D = 11 supergravity,” *JHEP* **09** (2006) 041, [arXiv:hep-th/0607093](#).